


Q11
.M76
15:4

Intermountain Journal of Sciences

(ISSN 1081-3519)

Vol. 15, No. 4 - 2009

JS



INTERMOUNTAIN JOURNAL OF SCIENCES

The Intermountain Journal of Sciences is a regional peer-reviewed journal that encourages scientists, educators and students to submit their research, management applications, or view-points concerning the sciences applicable to the intermountain region. Original manuscripts dealing with biological, environmental engineering, mathematical, molecular-cellular, pharmaceutical, physical and social sciences are welcome.

Co-sponsors/publishers include the Montana Academy of Sciences, the Montana Chapter of The Wildlife Society, and the Montana Chapter of The American Fisheries Society. This journal offers peer review and an opportunity to publish papers presented at annual meetings of the co-sponsor organizations. It is the intent of the governing bodies of the co-sponsor organizations that this journal replace printed proceedings of the respective annual meetings. Therefore, it is the policy of the editorial board that presenters at annual meetings of the co-sponsors be given priority in allocation of space and time of publication, although submission of other manuscripts for review and publication without regard to membership is encouraged.

Initial funding was provided by the co-sponsor organizations. Long-term funding will be derived from page charges assessed authors, sponsoring organizations or agencies at \$60 per printed page upon acceptance of each manuscript and from annual subscriptions: student \$6; regular member \$15; patron member \$25; overseas member \$25; library \$25; life member \$150; and, sustaining subscriber \$2,500.

The intent of the co-sponsors and editorial board is that *The Intermountain Journal of Sciences* be expanded to a quarterly journal. Achieving that objective depends upon numbers of acceptable manuscripts received and available funding. It also is the intent of the editorial board that contributing authors be assured of publication within 12 months of acceptance of their manuscript by the managing editor.

The organizational staff is voluntary and consists of an editorial board, an editor-in-chief, a managing editor, associate editors, a business manager and a panel of referees. The editorial board is responsible for establishing policy and the chair of the editorial board serves as liaison to the sponsoring organizations. The editor-in-chief is responsible for determining acceptability and level of revision of manuscripts based on referees' comments and recommendation of an associate editor. The managing editor serves as liaison for layout and printing. Associate editors include but are not limited to the section vice presidents of The Montana Academy of Sciences. Referees are selected on the basis of their field and specific area of knowledge and expertise.

Referees and associate editors judge submitted manuscripts on originality, technical accuracy, interpretation and contribution to the scientific literature. Format and style generally follow the *Guidelines for Manuscripts Submitted to the Intermountain Journal of Sciences, Dusek 1995, revised 2007*.^{*} Organization may vary to accommodate the content of the article, although the text is expected to elucidate application of results.

^{*}For detailed information about IJS, please go to our web site at:
www.intermountainjournal.org

INTERMOUNTAIN JOURNAL OF SCIENCES

EDITOR-IN-CHIEF

Gary L. Dusek, Spokane Valley, WA

MANAGING EDITOR

Terry N. Lonner, Bozeman, MT

ASSOCIATE EDITORS

BIOLOGICAL SCIENCES

Robert Harrington - Botany
Montana Fish, Wildlife and Parks
1400 S. 19th Avenue
Bozeman, MT 59718

David Stagliano - Aquatic Ecosystems
Montana Natural Heritage Program
1515 E. 6th Avenue
Helena, MT 59620-1800

Amy J. Kuenzi - Terrestrial Ecosystems
Department of Biology
Montana Tech of the Univ. of Montana
Butte, MT 59701

ENVIRONMENTAL SCIENCES AND ENGINEERING

Holly G. Peterson
Environmental Engineering Dept.
Montana Tech of the Univ. of Montana
Butte, MT 59701

HUMANITIES AND SOCIAL SCIENCES

Ismail H. Genc
College of Business and Economics
University of Idaho
Moscow, ID 83844

MATHEMATICS, STATISTICS AND COMPUTER SCIENCE

Keith Olson
Dept. of Computer Sciences
Montana Tech of the Univ. of Montana
Butte, MT 59701

MOLECULAR CELLULAR BIOLOGY AND NEUROSCIENCES

Richard Bridges
School of Pharmacy
University of Montana
Missoula, MT 59812

PHARMACOLOGY AND TOXICOLOGY

Charles Eyer
School of Pharmacy
University of Montana
Missoula, MT 59812

PHYSICAL SCIENCES

Richard Smith
Physics Department
Montana State University
Bozeman, MT 59717

EDITORIAL BOARD

Richard J. Douglass
Scott Barndt
James Barron
Robert G. Bramblett
Thomas Komberec
Carter Kruse
Tom Lewis
Mike Thompson
John P. Weigand

Chair, Montana Tech of the University of Montana - Butte
USDA Forest Service - Bozeman, MT
Montana State University - Billings
Montana State University - Billings
USDA Forest Service, Retired - Drummond, MT
Turner Enterprises, Inc. - Bozeman, MT
Montana State University - Billings
Montana Fish, Wildlife & Parks - Missoula
Montana Fish, Wildlife & Parks, Retired - Bozeman

BUSINESS MANAGER

Fred Nelson

Bozeman, MT

FINANCIAL STATEMENT (1/01/09 - 12/31/09)

Balance 01/01/09	\$508.65
Income:	
Regular Member	\$90.00
Library Subscriptions	\$300.00
International Member	\$25.00
Subscriptions Total	\$415.00
Page Charges	\$5,740.00
Reprints	\$100.00
Back Issue Sales	\$35.15
<hr/>	
Total Income	\$6,290.15
Expenses:	
Postage	\$330.69
Office Supplies	\$68.78
P. O. Box Rent	\$96.00
Misc. Fees	\$15.00
Photocopy	\$10.68
Total Expenses	\$521.15
 Balance 12/31/09	 \$6,277.65

Fred Nelson, Business Manager

EDITORIAL REVIEW POLICY

The *Intermountain Journal of Sciences* (IJS) is a fully refereed journal.

Manuscripts are submitted to the Editor-in-Chief (EIC) for initial consideration for publication in the IJS. This review shall include, but not be limited to, appropriateness for publication in this journal, correct formatting, and inclusion of a letter of submittal by the author with information about the manuscript as stated in the “Guidelines for manuscripts submitted to the *Intermountain Journal of Sciences*” (Dusek 1995, 2007). This cover letter must also include a statement by the author that this paper has not been submitted for publication or published elsewhere. The EIC notes the date of receipt of the manuscript and assigns it a reference number, IJS-xxxx. The EIC forwards a letter of manuscript receipt and the reference number to the corresponding author. The corresponding author is the author who signed the submittal letter.

Three hard copies of the submitted manuscript, with copies of the “Guidelines and checklist for IJS referees” attached are forwarded to the appropriate Associate Editor. The Associate Editor retains one copy of the manuscript and guidelines for his/her review, and submits a similar package to each of two other reviewers. A minimum of two reviewers, including the Associate Editor, is required for each manuscript. The two other reviewers are instructed to return the manuscript and their comments to the Associate Editor, who completes and returns to the EIC a blue “Cover Form” and all manuscript and reviewer comments plus a recommendation for publication, with or without revision, or rejection of the manuscript. This initial review process is limited to 30 days.

The EIC reviews the recommendation and all comments. The EIC then notifies the corresponding author of the results of the review and the publication decision.

ACCEPTANCE

For accepted manuscripts, each copy of the manuscript containing comments thereon and other comments are returned to the corresponding author. Revised manuscripts are to be returned to the EIC in hard copy, four copies if further review is required, or one hard copy plus the computer disk if only minor revision or formatting is necessary. The revised manuscript shall be returned to the EIC within 14 days of the notification. Review of the revised manuscript by the Associate Editor and reviewers shall be completed and returned to the EIC within 14 days. An accepted manuscript will then be forwarded to the Managing Editor (ME) for final processing.

REJECTION

Each manuscript that is rejected for publication is returned by the EIC to the corresponding author along with the reasons for rejection. The author is also advised that the manuscript may be resubmitted, provided all major criticisms and comments have been addressed in the new manuscript. The new manuscript may be returned to the initial review process if deemed appropriate by the EIC. If the manuscript is rejected a second time by either the EIC or the Associate Editor and reviewers, no further consideration will be given for publication of the manuscript in IJS. The corresponding author will be notified of this decision.

REVIEWER ANONYMITY

The identity of all reviewers shall remain anonymous to the authors, called a blind review process. All criticisms or comments by authors shall be directed to the EIC; they may be referred to the ME or the Editorial Board by the EIC for resolution.

MANUSCRIPTS SUBMITTED BY EDITORS

Each manuscript submitted by an Associate Editor shall be reviewed by the EIC and a minimum of two other reviewers with expertise in the subject being addressed. Each manuscript submitted by the EIC shall be forwarded with the necessary review materials to the Chairman of the Editorial Board of IJS, who will serve as the EIC for that manuscript.

ABSTRACTS

Only abstracts from the annual meetings of the sponsoring organizations will be published in IJS. Other submissions of abstracts shall be considered on a case-by-case basis by the Editorial Board. Sponsoring organizations shall collect abstracts, review them for subject accuracy, format them in Microsoft Word and email them to Gary Dusek, the EIC (gnpdusek@msn.com), on or before November 1. Each abstract shall be reviewed by the EIC to assure proper grammar, compliance with IJS "Guidelines for Abstracts Only" and

for assignment to the appropriate discipline section. All abstracts will be published in the December issue only.

COMMENTARY

Submissions concerning management applications or viewpoints concerning current scientific or social issues of interest to the Intermountain region will be considered for publication in the "Commentary" Section. This section will feature concise, well-written manuscripts limited to 1,500 words. Commentaries will be limited to one per issue.

Submissions will be peer reviewed and page charges will be calculated at the same rate as for regular articles.

LITERATURE CITED

Dusek, Gary L. 1995, revised 2007.

Guidelines for manuscripts submitted to the *Intermountain Journal of Sciences*. Int. J. Sci. 1(1):61-70. Revised guidelines are available on the Intermountain Journal of Sciences web site: (www.intermountainjournal.org)

EFFECTS OF A 6-WEEK JUMP TRAINING PROGRAM COMBINED WITH TWO LIFTING PROGRAMS

John Amtmann, Ed. D., Safety, Health and Industrial Hygiene Dept. Montana Tech, Butte, MT 50701

Jeremy Basler, Applied Health Science Student, Montana Tech, Butte, MT 50701

Charles S. Todd, Ph.D., Math and Statistics Department, Montana Tech, Butte, MT 50701

Shawn Vincent, Applied Health Science Student, Montana Tech, Butte, MT 50701

ABSTRACT

We evaluated the effects of a 6-week jump training program on 12 college basketball players using two strength training programs. The Leg Press (LP) group participated in a strength program using the leg press, and the Power Clean (PC) group used the power clean exercise instead of the leg press. Both groups participated in a 6-week jump training program. Pre-program measurements recorded jump scores for one-foot take off and two-foot take offs. The LP group had an average pre-program vertical of 66.68 and 88.05 cm for the 2-foot and 1-foot take offs, respectively. The PC group had an average pre-program vertical of 72.60 and 88.05 cm for the 2-foot and 1-foot take offs, respectively. The LP group had an average post-program vertical of 70.91 cm for the 2-foot take-off for an increase of about 4.2 cm, and 89.75 cm for the 1-foot take-off for an increase of about 1.7 cm. The PC group had an average post-program vertical of 76.2 cm for the 2-foot take-off for an increase of 3.6 cm, and 90.81 cm for the 1-foot take-off for an increase of about 2.75 cm. The comparison of average increases between the LP and PC groups was not significant ($P \geq 0.05$) for either the 2-foot or 1-foot take off.

Key words: jump training, vertical jump, power clean, leg press

INTRODUCTION

A common quest for athletes is to become more explosive or more powerful. The most dominant athletes in sports often are more powerful than their opponents. Two general theories prevail regarding how strength training improves performance. One theory says that strength training exercises that mimic sports skills may improve power that transfers onto the particular field of play (Fleck and Kraemer 2004). Strength training exercises that mimic sports skills are usually Olympic lifts or their derivatives such as the power clean. Strength training professionals that adhere to this theory believe that in order to become explosive, an athlete must train explosively in the weight room (Pulkkinen 2001, Lansky 1999, Fleck and Kraemer 2004, Grisaffi 1996, Takahishi 1992).

The second theory holds that no transfer of skills occurs between strength training movements and sport movements. Additionally, these theorists believe that

explosive exercises done in the weight room will significantly increase the likelihood of injury (Stein 1999, Mannie 1996, Brzycki 1995). Strength training professionals who embrace this theory believe two requirements are necessary for becoming more powerful and explosive within a specific sports arena. First, the athlete must practice specific movements with perfect technique, literally thousands of times so that the technique becomes automatic. Second, the athlete should strengthen all of the muscle groups, in a balanced fashion throughout a full range of motion. Increased power and explosiveness will occur as a result of improvements in overall strength and efficiency of movement involved in a specific skill. Supporters of this theory believe that sports movements should not be mimicked in the weight room because this practice may increase the chance of injury, and will do nothing to improve the sport-specific efficiency of movement (Brzycki 2002, Mannie 1999, Stein 1999).

Research Question/Hypotheses

With these theories in mind, we implemented a 6-wk jump training program for the men's basketball team at Montana Tech. Half the athletes participated in a strength training program that utilized the power clean, and the other half participated in a strength training program that used the leg press instead of the power clean; and all participants participated in a 6-wk jump training program. Is the power clean the preferred exercise for developing jumping power? To consider that question we developed the following hypotheses:

Null Hypothesis 1— a difference in the average increase (post-pre) of 2-foot take off jump scores between the leg press and power clean groups following a 6-wk jump training program is not statistically significant ($P \geq 0.05$).

Research Hypothesis 1— an increase in the average increase (post-pre) of 2-foot take off jump scores between the leg press and power clean groups following a 6-week jump training program is statistically significant ($P \leq 0.05$).

Null Hypothesis 2—a difference ($P \geq 0.05$) in the average increase (post-pre) of 1-foot take off jump scores between the leg press and power clean groups following a 6-wk jump training program is not statistically significant.

Research Hypothesis 2— an increase ($P \leq 0.05$) in the average increase (post-pre) of 1-foot take off jump scores between the leg press and power clean groups following a 6-wk jump training program is statistically significant.

METHODS

Subjects/Procedures

Twelve athletes ranging in age from 18-23 yrs from the Montana Tech men's basketball team volunteered for this study. Half of the athletes were assigned to the normal strength training program (power clean group) provided by the basketball coaches, and the other half of the athletes were assigned to the same program with the following change: the power clean

that is normally performed was replaced with the leg press (leg press group). So the athletes participated in the same strength training program except one group used the power clean and the other used the leg press. Exercises for the strength training programs included leg press/power clean, leg extension, leg curl, abdominal curl, hyperextension, bench press, lat pull down, triceps extension and bicep curl. The strength training program was combined with a jump training program based on the recommendations of Michigan State University's Strength Coach, Mike Vorkapich, in his article Jump Training (2001). Exercises included a warm-up and followed with various combinations of walking thigh pulls, jump skips, tuck jumps, frog jump, backboard slaps and x-out layups. We used a Vertec vertical leap measuring device to assess both sets of jump scores on an evening when the athletes had one day of rest from practice and strength training. Each athlete was allowed two jumps for each take-off, and the initial jump scores were compared with jump scores tallied at the end of training.

Both groups participated in a pre-program 2-foot static take off and 1-foot dynamic take off vertical jump test prior to a 6-wk jump training program. Jump training was a 2 day/wk program that implemented exercises specific to basketball jumping with both one- and two-foot take-off drills. The program consisted of a range of 80-110 foot jumps, leaps and bounds, and the duration of the program was approximately 20 min/session. Following the 6-wk jump-training, subjects completed the same vertical jump test to detect any improvement.

RESULTS

The LP group had an average pre-program vertical of 66.68 and 88.05 cm for the 2-foot and 1-foot take offs, respectively (Table 1). The PC group had an average pre-program vertical of 72.60 and 88.05 cm for the 2-foot and 1-foot take offs, respectively (Table 2). The LP group had an average post-program vertical jump of 70.91 cm for the 2-foot take-off for an increase of about

Table 1. Leg Press (LP) Group Results.

Leg Press	Beg 1 ft	End 1 ft	Beg 2 ft	End 2 ft
LP1	91.44	96.52	69.85	71.12
LP2	85.09	90.17	57.15	63.5
LP3	91.44	90.17	64.77	72.39
LP4	81.28	82.55	60.96	64.77
LP5	86.36	87.63	69.85	74.93
LP6	92.71	91.44	77.47	78.74
Avg	88.05	89.75[#]	66.68	70.91
		<i>P</i> = 0.4189		
			<i>P</i> = 0.6190	

Table 2. Power Clean (PC) Group.

Power Clean	Beg 1 ft	End 1 ft	Beg 2 ft	End 2 ft
PC 1	83.82	86.36	69.85	72.39
PC2	95.25	97.79	78.74	83.82
PC3	82.55	85.09	67.31	71.12
PC4	76.2	83.82	67.31	72.39
PC5	85.09	87.63	71.12	71.12
PC6	105.41	104.14	81.28	86.36
Avg	88.05	90.81	72.60	76.20

4.2 cm, and 89.75 cm for the 1-foot take-off for an increase of ~1.7 cm. The PC group had an average post-program vertical of 76.2 cm for the 2-foot take-off for an increase of 3.6 cm, and 90.81 cm for the 1-foot take-off for an increase of about 2.75 cm.

Due to the relatively small sample sizes ($n = 6$ for both the LP and PC groups) the non-parametric Mann-Whitney test was conducted to determine statistical significance between the groups. Both groups improved their average vertical jump performance, which we expected.

Results were not statistically significant when comparing the average increase for the 2-foot Jump between the LP and PC groups ($P > 0.05$). Thus, we failed to reject null hypothesis 1. Similar results were obtained when we compared the average increase between the two groups for the 1-foot jump ($P > 0.05$). That is, we also failed to reject null hypothesis 2.

DISCUSSION

Use of the power clean to improve athletic performance is an intensely debated topic. Results of this project allow us to conclude that the power clean was no more effective than the leg press, to improve vertical leap scores for these groups when combined with a jump-training program.

The leg press is a simple exercise that is safe to perform. The power clean is a complex exercise that has more danger factors as a result of significantly higher compressive and shear forces than the leg press with potential injuries occurring to the wrist, elbow, shoulder, back, hip and knee (Brzycki 2002, Kielbaso 1999).

Limitations to our study include the fact that our convenience sample was small and is not necessarily representative of the entire population. Also, we should have tracked their strength changes during the 6-wk period. We recommend that future research include tracking strength changes, using larger sample-sizes that include a control group, and conducting similar studies on females.

LITERATURE CITED

- Brzycki, M. 2002. *Wrestling strength: prepare to win*. Cardinal Publishers Group: Indianapolis, IN.
- Fleck, S. and W. Kraemer. 2004. *Designing resistance training programs*. Human Kinetics. Champaign, IL.
- Gill, N. 2002. Email interview. September, 2.
- Grisaffi, D. 1996. Ballarmine Prep's Strength Training Program for Wrestling. *Strength and conditioning* 18: 54-58.

- Kielbaso, J. 1999. The Reality of the Power Clean. *In* M. Brzycki, editor, Maximize your training: insights from leading strength and fitness professionals. Masters Press, Chicago, IL.
- Lansky, R. 1999. Wrestling and Olympic Style Lifts: In-Season Maintenance of Power and Anaerobic Endurance. *Strength and Conditioning Journal* 21:21-27.
- Mannie, K. 1999. Athletic Skill Development: An Open and Closed Case. *In* M. Brzycki, editor, Maximize your training: Insights from leading strength and fitness professionals. Masters Press, Chicago, IL.
- Pedro, J. 2002. Phone interview. September, 4.
- Pulkkinen, W. 2001. The sport science of elite judo athletes. *Pulkinetics*, Guelph, Ontario, Canada.
- Stein, A. 1999. Improving Athletic Power. *Hard Training* Issue 8: 7-9.
- Takahashi, R. 1992. Power Training for Judo: Plyometric Training with Medicine Balls. *National Strength and Conditioning Association Journal* 14: 66-71.
- Vorkapich, M. 2001. Jump Training. Unpublished manuscript.

Received 18 May 2009

Accepted 18 December 2009

ABSTRACTS

THE MONTANA CHAPTER OF THE AMERICAN FISHERIES SOCIETY
AND

THE MONTANA CHAPTER OF THE WILDLIFE SOCIETY

2009 JOINT CONFERENCE

ADAPTING FISH AND WILDLIFE MANAGEMENT TO HUMAN DEMOGRAPHIC CHANGE IN MONTANA

FEBRUARY 9 – 13, 2009

KALISPELL, MONTANA

Nearly 400 members of the Montana Chapters of the American Fisheries Society and The Wildlife Society gathered in Kalispell, Montana, February 9-13, 2009 for a joint meeting. The meeting's theme was "*Adapting Fish and Wildlife Management to Human Demographic Change in Montana*", recognizing that successfully solving looming fish and wildlife management challenges such as climate change and expanding energy development can only occur when human demography is included in solutions. The meeting began with two continuing education workshops designed to provide both scope and solutions to elements of the theme: *Human Demographic Impacts: Invasive Species and Other Species of Concern*, and *Human Demographic Impacts: Solutions to Fish and Wildlife Management Challenges*. Then, the plenary session, titled *Learning from the past, but focusing on the future – adapting fish and wildlife management to Montana's new demographics* presented both the progression of fish and wildlife management in Montana in response to human demographic changes over the last century, as well as a context for future changes. The plenary session concluded with panel discussions that provided perspectives from both public and private entities spanning most aspects of fish and wildlife management. A consistent message to the audience was that human demographic change is modifying some fundamental tenets of fish and wildlife management, and therefore will require development of both new models of management and collaboration, and different skill sets among future managers. Although this change presents challenges, it likewise offers exciting opportunities for disparate disciplines and groups to work together to solve these challenges. As such, it was very appropriate that the topic was addressed by a joint meeting of the two largest professional fish and wildlife societies in Montana.

After the plenary session, a day of three joint concurrent sessions ensued. Presentations were grouped in topics of mutual concern to both fish and wildlife managers. Consequently, both topics and attendees were diverse and interdisciplinary. The abstracts of those presentations at the 2009 joint meeting demonstrate the wide reaching, important and timely work conducted by Montana's fish and wildlife professionals. Based on this and past examples, there is no doubt that the dedicated individuals, agencies, and organizations represented by the Montana Chapters of The Wildlife Society and the American Fisheries Society will continue to strive to meet the challenges posed by a changing future.

Scott Barndt, President, Montana Chapter of the American Fisheries Society
Joe Weigand, President, Montana Chapter of The Wildlife Society

JOINT PLENARY INVITED ABSTRACTS

MONTANA'S WILDLIFE LEGACY – DECIMATION TO RESTORATION

Harold Picton, Emeritus Professor of Wildlife Management, Department of Ecology, Montana State University, Bozeman, Montana 59717

Terry Lonner, Chief of Wildlife Research - Montana Fish, Wildlife and Parks (retired); Media Works - Multimedia Producer, Bozeman, Montana 59715

The main intent of the Montana Wildlife Legacy Project was to pay tribute to the generations of Montanans who made our current wildlife resources possible and for public education, especially in junior high and high schools and hunter safety classes. Use of the wildlife resources for the settlement of Montana produced a catastrophe by the end of the 19th century, but by the end of the 20th century wildlife was more abundant than at any time during the previous 130 years. Passage of protective laws during the latter 19th century, coupled with gradually increasing efforts to enforce those laws, accounted for a portion of this resource restoration. The remainder of the story is largely untold and is the subject of the DVD, "Back from the Brink - Montana's Wildlife Legacy" produced in 2005 and the companion book, "Montana's Wildlife Legacy - Decimation to Restoration" published in 2008. Game bird resources were heavily affected by the "cow and the plow." Few agricultural lands provided new habitats unsuited for native species. Pheasant introductions began before 1895 by private individuals and became a state bird farm program in 1929. Over 0.80 million pheasants were planted by the time the state discontinued this program in the early 1980s. At the beginning of the 20th century elk occurred only in the Sun River-South Fork of the Flathead and Yellowstone National Park areas. Rod and Gun clubs held fund-raising events and paid \$5/ elk to have the Northern Pacific Railroad deliver rail car loads of elk trapped near Gardiner, Montana just north of Yellowstone National Park, to areas near their towns. The transplants began in 1910 with releases near Butte, Hamilton and in the Glacier Park area. Transplants of elk continued until 1997 when the last of 11,364 were trapped and released. Similar programs for other species were developed after the Pitman-Robertson funds were accepted by the Montana Legislature in 1939 with the Wildlife Restoration Division of the Montana State Fish and Game Department established in 1940. For example, almost 4000 pronghorn antelope were trapped and transplanted, 1000 more than existed in the entire state in the 1920s. Significant, but regulated hunter harvests of the biological surplus and habitat preservation programs became prominent in the last third of the 20th century. This resurrection of Montana's wildlife resource was an epic effort extending through six generations, five wars, an economic collapse, and the greatest North American climate disaster of the 20th century. Citizen leadership arose in generation after generation and melded with the leadership and science furnished by the state, federal agencies, and universities to bring about a successful restoration effort. Wildlife resources that we enjoy today were a gift of the people from the 20th century to the people of the 21st century. They come with a message to care for and cherish them, to value and maintain them. Resources that are not valued tend to end up on the trash heaps of human history. For more information go to: www.montanawildlifelegacy.com.

CHRIS HUNTER'S INTERPRETATION OF HUGH ZACKHEIM'S HISTORY OF MONTANA FISH, WILDLIFE AND PARKS FISHERIES DIVISION 1901-2005

Chris Hunter, Montana Fish, Wildlife and Parks, P.O. Box 200701, Helena, Montana 59620

The Fisheries Division was created in 1901. The first resident licenses were created at the same time, with more than 30,000 of the \$1 licenses purchased in the first year. For the first 50 years the emphasis of the division was on stocking fish. The first fisheries biologist was hired in 1947. The results of scientific investigations by these biologists led to changes in fish management including changes in stocking policy and increased emphasis on the effect of environmental damage to Montana's fishery resources. A spate of new environmental laws in the 1960s and 1970s made significant progress toward protecting Montana's aquatic environments. In 1974 the Fish and Game Commission adopted the wild trout policy. The 1980s and 90s saw increasing emphasis on native fish, T&E species issues and continuing efforts to secure water for instream flows. There was also an increased emphasis on habitat restoration with the Future Fisheries Program, hydro relicensing and Natural Resources Damages suits. These areas of emphasis have continued into the new millennium. State and national trends for hunting and fishing are clearly on the decline. There is increasing concern about the lack of time our children spend involved in nature related activities. What are the implications for the future of our agencies and the resources we manage?

THE MONTANA CHALLENGE

SuzAnne M. Miller, Dunrovin Research, P.O. Box 822, 5375 Terry Lane, Lolo, Montana 59847

The Montana Challenge begins with a question posed by the USDA Forest Service and Montana Fish, Wildlife and Parks. How can management of Montana's fish and wildlife resources best contribute to Montana's social and economic well being? By employing prominent socioeconomic researchers from across the Rocky Mountain region, and digging for authoritative public-sector data, The Montana Challenge established an unexpectedly vital role of clean air, clean water, fish, wildlife, and wild lands in Montana's culture and economy. The Rocky Mountains are the fastest growing section of the U.S. in population, personal income, and total employment; while the Great Plains are in population and economic decline. Montana's counties reflect this trend with rapid growth along the mountains in the west, slower growth along the Rocky Mountain front, and rapid decline in the eastern plains. "Quality of life" is driving the Rocky Mountain West's population and economic growth and natural resource amenities are key to quality of life and economic prosperity. Montana and other Rocky Mountain states are in the midst of a transition from an economy based on natural resource commodities to a human resource based economy where jobs follow the people. Montana's prosperity depends on attracting people who create economic opportunities. Healthy ecosystems, healthy fish and wildlife populations, and broad public access create Montana's unique and desirable lifestyles. Montana's wild resources draw people and economic activity. Diversity is a source of strength for both ecosystems and economies. Human resource sectors are the most rapidly growing sectors of Montana's economy, especially in the western region. Montana's traditional natural resource based economic activities remain important to Montana, especially in the eastern region. Montana's challenge is to both utilize and protect its natural resources.

FISH AND WILDLIFE VALUES IN THE WEST

Michael J. Manfredo and Tara L. Teel, Human Dimension of Natural Resources, Department, Colorado State University, Fort Collins, Colorado

Western states are going through a number of changes that have affected and will continue to affect wildlife management. Changes include population growth, changes in in-migration rates and land ownership patterns, increasing income and education levels, growth in technology, and urbanization. The recent study, **Wildlife Values in the West**, explores how some of these broad societal forces are shaping the composition of public values toward wildlife throughout the western region. **Wildlife Values in the West** is a project of the Western Association of Fish and Wildlife Agencies Human Dimensions Committee. It is a collaborative regional effort involving social science researchers from Colorado State University and representatives from 19 participating state fish and wildlife agencies who assisted in development of a mail survey. Data were collected through administration of the survey to a sample of residents in each state in the fall of 2004 ($n = 12,673$). Results and related implications from this 19-state effort will be discussed, including the impacts value shift may have on public acceptance of wildlife management strategies and on demand for participation in wildlife-related recreation activities in Montana. We will also discuss how study results provide a broad context to assist state fish and wildlife agencies better understand diverse publics and plan for the future of wildlife management in the west.

GREATER YELLOWSTONE AREA ELK MOVEMENTS: BRUCELLOSIS RISK AND HUNTER ACCESS

Kenneth L. Hamlin and Julie A. Cunningham, Montana Fish, Wildlife and Parks, 1400 S. 19th Ave., Bozeman, Montana 59718

Thomas O. Lemke, Montana Fish, Wildlife and Parks, 406 Chestnut Lane N. Livingston, Montana 59047

We examined elk movement data from across the Greater Yellowstone Area (GYA) from 1976-2006, comparing and contrasting movement patterns within and between herds. Our objectives were to understand how public hunter access related to elk movements and brucellosis risk in different areas of the GYA. We focused on the Madison Valley, Gallatin Valley and east side Paradise Valley. In the Madison Valley, we compared elk movement dynamics between 27 cow elk monitored 1976-1986 (VHF collars) and 43 cow elk monitored 2005-2006 (GPS collars). Over this time period, land ownership changes resulted in reduced hunter access to private lands for cow elk hunting. We found that, compared to the 1976-1986 movements, 2005-2006 elk migrated earlier to wintering ranges, left later to summer ranges, and used private land areas more extensively. During 2005-2006, cow elk were less available to hunters due to use of private land refuges during the hunting season. In the Northern Yellowstone, we compared elk movements from 1984-1987 (VHF collars) with preliminary data from 2007-2008 (GPS collars). Preliminary analysis suggests that elk migrated to and from winter range generally with expectation given weather conditions. Some individuals were more available to public hunters than others (range = 0-100%) based on behavioral and movement patterns. Flight and GPS data from 2007-2008 elk indicate Northern Yellowstone elk spend summertime further south and west in Yellowstone National Park than had previously been thought, and that Northern Yellowstone elk may be coming into contact with Jackson Hole, WY, elk on these ranges.

A FISHERY MANAGER'S CHALLENGE

Jim Vashro, Montana Fish, Wildlife and Parks, 490 N. Meridian, Kalispell, Montana 59901

Fisheries management is increasingly defined by changes in human populations, loss of access to public waters, native fish management and the struggle to maintain aquatic habitat in the face of both climate change and physical alterations. Angling pressure has not kept pace with population growth and combined with the decline of rod and gun clubs and the increase in special interest groups threatens both funding and social and political support for aquatic resource management. Changes in land-ownership funnels that increases numbers through fewer access sites results in conflicts. Changing land ownership also leads to increasing applications and violations for fish ponds and streambank alterations and old financial constraints do not always apply. Many anglers do not have a good grasp of biological principles that leads to illegal fish introductions and challenges to management programs. Today's biologists are better trained than ever but recruitment and retention is an increasing problem for Fish, Wildlife and Parks. Many management tools have not changed but new technologies offers greater understanding if we use them intelligently. Native fish management increasingly drives management through statutes, policies and funding priorities. Suppression of unwanted species is the latest strategy but not always applied well. The angling public has not been educated well on the need for native fish management and often challenges programs. Climate change and a growing demand for energy development could trump many of today's programs.

HUNTER ACCESS – A MOVING TARGET FOR MANAGERS

Alan Charles, Montana Fish, Wildlife and Parks, P.O. Box 200701, Helena, Montana 59620-0701

Traditionally, fish and game agency program managers deal with science as biologists or laws as game wardens. Agencies typically adopt regulations, issue licenses, and hunters set forth on their own to find a place to hunt. However, in recent years, Montana Fish, Wildlife, and Parks has found it necessary to develop access programs and become more involved in helping ensure that the hunters who buy the licenses have a place to hunt. This has become necessary not only to maintain Montana's hunting heritage and traditions, but also to ensure that public hunting remains an effective tool to help manage the state's populations of deer, elk, and antelope. Changes in who owns the land, how the land is managed, what hunters expect, and what hunters are willing to do all pose challenges for wildlife agency program managers as the world changes around them.

PROTECTING PUBLIC RESOURCES ON PRIVATE LAND: ADAPTING BUSINESS STRATEGIES TO CHANGING DEMOGRAPHICS AND DEMANDS

David Greer, Plum Creek Timber Company, Inc. Columbia Falls, Montana 59912

Plum Creek is the largest private forest landowner in the United States with 7.4 million acres in 19 states and 1.1 million acres in Montana. While Plum Creek's core business is timber management, real estate transactions have always been a part of the business, including conservation transactions and land sales to private entities. Conservation strategies, beyond easements and sales, are integrated into routine management efforts including adherence to the standards of the Sustainable Forestry Initiative ®, habitat conservation plans, cooperative

agreements, and land exchanges. Since 1989, conservation transactions to public agencies, conservation organizations, and timber companies have accounted for 81 percent of all Plum Creek land sales in Montana. Additionally, conservation easements sold to public agencies in Montana have amounted to more than 149,000 acres and include one of the largest easements completed in the U.S. Private parties seeking Montana property increasingly want to be viewed as “green” and consequently are more receptive to deed restrictions and protection practices that address public resources, such as grizzly bears and native fish habitat. The “recipe” for success in conservation land transactions includes (1) availability of large, strategically important tracts, (2) willingness by Plum Creek to work with innovative partnerships, and (3) patience; and adherence to a “win-win” strategy for the company and the public. The potential for future conservation transactions in Montana will hinge on finding creative financing solutions and incorporating active forest management provisions to support local timber-based economies. The Montana Working Forest Project, seeking to transfer 320,000 acres of Plum Creek property to federal, state, and other private ownership is an example of this type of conservation transaction.

PERSPECTIVE OF A NON-TRADITIONAL LANDOWNER/MANAGER

Russ Miller, Turner Enterprises, Inc. Bozeman, Montana 59718

TEI’s philosophy is to manage their lands in an economically sustainable and ecologically sensitive manner, while promoting the conservation of native species.

MONTANA’S LANDOWNERS: MANAGEMENT IMPLICATIONS OF MONTANA’S CHANGING DEMOGRAPHICS FOR WATER RIGHTS

Stan Bradshaw, Trout Unlimited, Montana Water Project, P.O. Box 412, Helena, Montana 59624

Water rights, long recognized as a property right in Montana, are among the least understood of all property rights. It’s not like a chunk of land—it’s a “use right.” A viable water right doesn’t exist simply as a statement on a certificate. If it is not put to a beneficial use, it can be lost. For over 100 years, it was widely believed that water had to be diverted to establish a water right. Water left in stream was a waste. Over the past four decades, Montana’s changing demographic—a gradual shift to a less rural population--first enabled the passage of instream flow legislation in the 1960s and 1970s and then again in the 1980s and 1990s when the legislature authorized leasing. So the first management implication is that we now have some tools. The changing demographic doesn’t always understand the limitations of those tools, but we have tools nonetheless. The other changing demographic is the influx of a new species onto traditional ranch lands—the amenity buyer. Always well-heeled, often well-intentioned, they can occasionally cause more problems than they can solve. On the other hand, the newcomers bring with them a different perspective that can enhance watershed restoration in ways not previously possible. The greatest challenge, will be educating both the traditional landowner and newcomer alike in the limitations of their water rights, while showing them the possibilities of creative change to those rights.

CHANGING HUMAN DEMOGRAPHICS INFLUENCE FOREST SERVICE MANAGEMENT IN MONTANA

Jane Cottrell, USDA Forest Service, Northern Region, P.O. Box 7669, Missoula, Montana 59807

Human demographic changes in Montana have been occurring for some time, bringing both challenges and opportunities for Forest Service Managers. Movement of people into urban/forest interfaces have created social, political and legal interest and action regarding the management of forests with respect to both fire and wildlife habitat. Human demographic changes have brought increasingly diverse perspectives to bear on issues surrounding management of the National Forests, making analysis and decision on issues such as travel management very complex. Emerging issues involving open space, aquatic health, invasive species and instream flow needs have also shaped management on the National Forests in Montana. We have also seen recent successes where diverse public interests have come together to resolve issues.

STREAM ACCESS: PAST, PRESENT AND FUTURE

Robert N. Lane, Montana Fish, Wildlife and Parks, Helena, Montana 59620-0701

The public's right to recreate in streams, rivers, and lakes will be discussed starting with its past roots in public trust concepts, covering how the present statutes and case law define the public use of streams and rivers, and concluding with an examination of how ongoing and future controversies will or may continue to shape rights of the public to recreate in streams, rivers, and lakes. The presenter covered (1) how Montana's Stream Access Law and rules and the Natural Streambed and Land Preservation Act (SB 310 Law) protect and help direct management of streams and rivers, (2) how these laws and rules are balanced with private property rights, and (3) what the public trust means for fisheries managers.

THE FUTURE AIN'T WHAT IT USED TO BE!

Bruce Farling, Montana Trout Unlimited, 240 North Higgins, Suite 12, Missoula, MT 59802

Montana's demographic landscape is comprised of three states within a state: The rapidly growing West, the modestly growing Central and the shrinking East. Close examination of each reveals important value shifts affecting fish, wildlife, recreation and habitat protection. Among these shifts are reduced opportunities for access to private lands for hunting and fishing, producing either a corresponding increase in pressure on public lands or fewer people choosing to hunt and fish. In response, managers will have to re-evaluate geographic and demographic priorities. The state's new demographic profile, which includes counties with high population growth rates that also have high rates of population turnover, means that traditional perspectives about wildlife and fish are in flux and influenced to an inordinate degree by values honed outside Montana or in an increasingly media-influenced world. Change elsewhere in the country, including that precipitated by a shifting climate, new resource demands, immigration, wealth transfer and the desire to live where the quality of life is high, could overwhelm the traditional bottlenecks that have moderated population growth in Montana, including winter weather, limited employment opportunities, and shortcomings in communication and transportation infrastructure. Demographic trends in Montana will necessitate new types of fish and wildlife managers, with different skill sets, tools and priorities that focus more on effective communication with greater attention to ecosystem protection, non-consumptive wildlife and fish values and equitable access to public resources possibly resulting in less attention to traditional management that focuses on hunters and anglers as primary beneficiaries.

BIOLOGICAL SCIENCES – AQUATIC

RURAL STAKEHOLDERS AND ARCTIC GRAYLING MANAGEMENT IN THE BIG HOLE RIVER WATERSHED, MONTANA, USA

Michelle L. Anderson, Kylene Owens, Biology Department, Montana Tech of the University of Montana, 1300 West Park St., Butte, Montana 59701

Michael A. Bias, Big Hole River Foundation, PO Box 3894, Butte, Montana 59702

In order to counteract Arctic grayling (*Thymallus arcticus*) population declines in the Big Hole River, rural stakeholders have partnered with natural resource managers on Candidate Conservation Agreements with Assurances (CCAA) activities. Surveys designed to assess attitudes towards grayling management practices were sent to 300 watershed residents in February 2008. We received 83 responses, mostly from men ≥ 50 years who identified their occupation as farming, ranching, government or retired. Respondents indicated grayling numbers had declined (31%) or stayed about the same (25%) in the last 10 yrs, and that it would be favorable for grayling numbers to increase (60%). Respondents chose drought, habitat loss and birds as factors strongly associated with declining grayling numbers. Activities listed as strongly associated with increasing grayling numbers varied, but often included drought management. Demographic trends among area residents match those typical of rural western communities; 62 percent are 45 years or older, 17 percent live below the poverty line, and 76 percent lack a college degree. The benefit of CCAA activities to the local “restoration” economy included an influx of nearly \$2 million since 2006. CCAA activities could be linked to future federal infrastructure, education and workforce training programs, with substantial benefits to the local populace.

DIVERSION DAMS AND FISH PASSAGE – DON’T GIVE UP THE FIGHT!

Mike Backes, Montana Fish, Wildlife and Parks, P.O. Box 1630, Miles City, Montana 59301

The Tongue River, in southeastern Montana, is a major tributary to the Yellowstone River. Numerous Yellowstone River fish species utilize the Tongue River for spawning. However, diversion dams have limited fish migrations up this system. Through persistent efforts, measured in careers not years, fish passage around these diversion dams and complete removal of other dams is occurring. T&Y Diversion dam, constructed in the 1880s, is the first dam migrating fish encounter, and it is a complete fish barrier. During fall 2007, a fish bypass (named Muggli Bypass) was completed to allow fish passage around T&Y Diversion Dam. Fish sampling was conducted in 2008 to evaluate the success of this structure. A fyke net was utilized to sample fish that successfully navigated the complete length of the bypass channel. Electrofishing was conducted upstream and downstream of the diversion dam to compare relative abundances of fish in the river to those collected in the bypass. Nineteen fish species were successful in passing through the bypass during the sampling period and four additional species were collected in the bypass channel when it was block netted and drained. Comparatively, 29 fish species were collected downstream of the diversion dam. Electrofishing upstream of T&Y Dam found four species (freshwater drum, goldeye, smallmouth buffalo and western silvery minnow) which have never been documented upstream of the dam. In summary, the Muggli Bypass is a noteworthy success story.

COMPETITION AS A FACTOR IN DISPLACEMENT OF NATIVE CUT THROAT TROUT BY NONNATIVE RAINBOW AND HYBRID TROUT

Laurie Battle, Department of Mathematical Sciences, Montana Tech, 1300 W. Park St., Butte, Montana 59701

Robert Van Kirk, Humboldt State University, 1 Harpst St., Arcata, California 95521

Bill Schrader, Idaho Department of Fish and Game, 1414 E. Locust Lane, Nampa, Idaho 83686

Native salmonid fishes have been displaced worldwide by nonnatives through competition and hybridization, but dynamics of these factors are poorly understood. We apply a Lotka-Volterra population model to displacement of cutthroat trout (*Oncorhynchus clarkii*) by rainbow (*Oncorhynchus mykiss*)/hybrid trout in the Snake River, USA. Cutthroat trout are susceptible to hybridization in the river but are reproductively isolated in tributaries via removal of migratory rainbow/hybrid spawners at weirs. Hybridization is the primary mechanism for initial growth of the rainbow/hybrid trout population, but a model with hybridization alone does not explain observed trends. Two models, in which competition occurs 1) among river-spawned fish only and 2) among all fish, explain observed trends, but are indistinguishable from one another based on fit to data. When tributary-spawned cutthroat trout out-migrate as fry, competition with rainbow/hybrid trout results in extinction of cutthroat trout, even though reproductive segregation is maintained.

PERSPECTIVES ON NOT LISTING THE FLUVIAL ARCTIC GRAYLING

Michael A. Bias, Big Hole River Foundation, P.O. Box 3894, Butte, Montana 59702

I summarize views held by the Big Hole River Foundation and provide insights into potential positive and negative outcomes of listing the fluvial Arctic grayling (*Thymallus arcticus*) under the Endangered Species Act. The perception that listing would lead to recovery of the grayling was evaluated based on past recovery of listed species. Currently, 1,925 species are on the Endangered Species List as either Threatened or Endangered. The number of species listed increased during the 1990s, but has decreased in recent years. To date, 48 species have been removed from the list; 22 have been "recovered," 17 have been reclassified due to data errors, and nine have gone extinct. For species in the contiguous U.S., 15 have been recovered, seven have gone extinct, and 15 have been removed due to data errors. The rate of recovery is estimated at 1.1 percent for North American species, and no fish species have been recovered to date. A common perception that a listing would increase funding was evaluated and showed that relatively little funds are available for species recovery under Section 6 of the ESA, when compared to other federal funding sources. Although the Foundation thinks evidence suggests a listing is warranted, they are supporting the CCAA as an appropriate means of addressing species recovery in the Big Hole.

TOXICITY OF ROTENONE TO LARVAL AMPHIBIANS

Hilary G. Billman, Sophie St. Hilaire, and Charles R. Peterson, Department of Biological Sciences, Idaho State University, Pocatello, Idaho 83209

Carter G. Kruse, Turner Enterprises, Inc., 1123 Research Drive, Bozeman, Montana 59718

Piscicide use in fisheries management is becoming increasingly common. Rotenone, specifically, is being used to remove non-native fish species from aquatic systems. While

the effects of this chemical on fish are well-studied, the impacts of rotenone on non-target species, such as amphibians, are not well known. This study was conducted to determine the toxicity of rotenone on two native species of tadpoles in Montana – *Rana luteiventris* and *Bufo boreas* – under laboratory conditions. For each species, tadpoles at three developmental stages were exposed to either a control or CFT Legumine (5% active rotenone) at one of four doses (0.1 mg/L, 0.5 mg/L, 1 mg/L, 2 mg/L). Total exposure time was 96 hours. Parameters measured included mortality at 96 hrs post treatment, and in the survivors, weight, Snout-Urostyle Length, and time to metamorphosis. In addition to the rotenone exposure trials, a recovery trial was conducted with early stage spotted frog tadpoles to determine survivability of tadpoles exposed to rotenone at 1 mg/L and then placed in clean water. Spotted frog tadpoles exposed to rotenone at 1mg/L – typical field application dose–experienced significantly greater mortality than control tadpoles. Although all stages of frog tadpoles exposed to rotenone were negatively affected by the chemical, the effect was worse at earlier life stages. Early stage toad tadpoles were significantly more resistant to rotenone exposure at 1 mg/L than early stage spotted frog tadpoles; however they were still negatively affected by the chemical. Sub-lethal effects, though statistically different between control and exposed survivors in two instances, were not consistent and therefore thought not to be biologically significant. Spotted frog tadpoles exposed to rotenone and then transferred to clean water experienced significantly lower mortality than those exposed for the full 96 hrs. Overall, rotenone exposure was found to be lethal to tadpoles of both species at all three developmental stages though mortality was not uniform across dosages or age groups.

BULL TROUT MONITORING; LOOK DEEP AND WIDE

Chris Clancy, Montana Fish, Wildlife and Parks, 1801 North First St., Hamilton, Montana 59840

Mike Jakober, USDA Forest Service, Bitterroot National Forest, West Fork Ranger District, 6335 West Fork Road, Darby, Montana 59829

In most of the Bitterroot drainage fluvial westslope cutthroat (*Oncorhynchus clarkii lewisi*) trout are fairly common and fluvial bull trout (*Salvelinus confluentus*) are rare. Monitoring the fluvial cutthroats is not difficult because mainstem population estimates are possible. However, due to the small number of fluvial bull trout, population estimates are not obtainable and redd counts are inconclusive. Monitoring of juvenile and resident populations also presents challenges. Population estimates (intensive) of bull trout that have the potential of providing quantitative data can be difficult to collect and basin-wide distribution (extensive) generally does not provide quantitative information. On the Bitterroot National Forest we have monitored bull trout and westslope cutthroat trout using mark-recapture population estimates for nearly 20 years. While these data allow us to follow trends within the reaches of a stream where populations are densest, it is not designed to identify trends in distribution. We have collected more extensive data but not in a systematic manner. We propose formalizing the collection of single pass electrofishing on short reaches throughout selected streams to augment the monitoring reaches established many years ago.

POPULATION CHARACTERISTICS OF LAKE TROUT IN SWAN LAKE, MT

Benjamin S. Cox and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Montana State University, Bozeman, Montana 59717

Wade A. Fredenberg, USDI Fish and Wildlife Service, 780 Creston Hatchery Rd., Kalispell, Montana 59901

Leo R. Rosenthal, Montana Fish, Wildlife and Parks, 490 North Meridian Rd., Kalispell, Montana 59901

The recent establishment of lake trout (*Salvelinus namaycush*) in Swan Lake, Montana threatens one of the most productive bull trout (*Salvelinus confluentus*) fisheries remaining in the USA. Management of invasive lake trout in other systems has been focused on suppression often without establishing a thorough baseline from which to evaluate the impacts of suppression efforts. Describing the population dynamics of lake trout in Swan Lake prior to suppression efforts will provide a baseline for evaluating the effects of exploitation in the future. In 2007 and 2008 extensive gill net sampling provided data to estimate size structure, density, condition, maturity, fecundity, age structure and mortality of lake trout. Size structure indices in 2007 and 2008 were low with proportional size distributions (PSD) values of 7 and 8 respectively. Lake trout density was estimated at 8800 (7300-10,500 95% CI) fish > 160 mm. Condition of lake trout in Swan Lake is among the highest recorded for populations in the northwestern USA. Relative weight (W_t) values varied from 92 for fish between 300 and 499 mm to > 120 for fish between 800 and 999 mm. Fifty percent of male lake trout mature at 584 mm and 50 percent of females at 726 mm. Fishing mortality was estimated between 36 percent and 52 percent based on the population estimate in 2008 and those fish removed. The thorough baseline established by this study provides managers with a reference point for evaluating the effects of exploitation on the lake trout population in Swan Lake, Montana.

CUTTHROAT TROUT STRUGGLE FOR PERSISTENCE IN THE FACE OF AN EXPANDING LAKE TROUT POPULATION IN YELLOWSTONE LAKE

Philip D. Doepke, Patricia E. Bigelow, Brian D. Ertel and Todd M. Koel, Center for Resources, Fisheries and Aquatic Sciences Program, P.O. Box 168, Yellowstone National Park, Wyoming 82190

Lake trout *Salvelinus namaycush* were discovered in Yellowstone Lake in 1994. The ecological threat of this non-native fish was immediately recognized and efforts to control lake trout were initiated the following year. The objective of the removal program is to reduce lake trout to the point where their effect on native Yellowstone cutthroat trout (*Oncorhynchus clarkii bouvieri*) is minimized. Lake trout capture techniques have focused on mechanical methods, mostly gill netting, and more recently electrofishing. Despite removal of almost 350,000 lake trout since 1995, with 76,140 removed in 2008 alone, all indicators point toward continued population growth. Catch per effort of juvenile lake trout has steadily increased since 2002 with 2008 being the highest on record (5.0 lake trout/100 m gill net set per night) since 1998. However, cutthroat trout have not shown a positive response to lake trout removal efforts. Numbers of spawning cutthroat trout in Clear Creek, a tributary to Yellowstone Lake, are at the lowest levels ever recorded and the lakewide cutthroat trout assessment catches are averaging 40 percent fewer cutthroat trout over the past 5 yrs when compared to pre-lake trout years. In 2008 a panel of experts convened to assess the current status of cutthroat trout and review lake trout removal efforts in Yellowstone Lake. The panel concluded that while current

levels of suppression have slowed lake trout population growth, more is required if a healthy cutthroat trout population is expected to persist in Yellowstone Lake. They concluded it was imperative that we immediately increase suppression efforts, and develop/implement lake trout monitoring.

ESTIMATING LAKE SURVIVAL OF JUVENILE BULL TROUT IN TRESTLE CREEK, IDAHO IN THE PRESENCE OF CHANGING FISH COMMUNITIES, LAND USE, AND FISH MANAGEMENT

Christopher C. Downs, Glacier National Park, Science Center, West Glacier, Montana 59936

Rob Jakubowski, Avista Corporation, PO Box 1469, Noxon, Montana 59853

Rob Ryan, Idaho Department of Fish and Game, 2885 W. Kathleen Ave., Coeur d'Alene, ID 83815

A total of 921 age-1 and older juvenile bull trout (*Salvelinus confluentus*) were marked emigrating from Trestle Creek, Idaho, from 2000 through 2002. Individual juvenile bull trout were marked with abdominally implanted PIT tags, and adult returns were monitored at an automated PIT tag detection weir, as well as at fish traps through 2008. All marked juveniles returned as adults by the end of the 2007 field season. Minimum estimates of survival from outmigrating juvenile to returning adult were similar across study years, and ranged from 8.9 percent to 15.5 percent. Short-term tag retention and survival of marked juveniles was high, but long-term tag loss estimated by examination of double-marked returning adult bull trout indicated substantial tag loss over time. Generally, outmigrating juveniles reared in the lake environment for between 3 and 5 growing seasons before returning as adults, with most spending four growing seasons in the lake (not including their "return year" as a year at-large in the lake). In general, return rates for larger outmigrants were higher than those for smaller outmigrants. Additional studies quantifying lake/river survival of migratory bull trout in other systems are needed to put these results into an appropriate ecological context, and to better understand the complex and likely interacting effects of non-native fish, land use, and fish management on bull trout recruitment.

BULL TROUT ENTRAINMENT AT LIBBY DAM IN THE KOOTENAI RIVER, MONTANA

Jim Dunnigan, Montana, Fish, Wildlife and Parks, 385 Fish Hatchery Road, Libby, Montana 59923

Montana Fish, Wildlife and Parks has used nighttime jetboat electrofishing to conduct annual mark-recapture population estimates of adult bull trout (*Salvelinus confluentus*) in the Libby Dam tailrace since 2004. Estimates are conducted during April or May within this 3.5-mi section of the Kootenai River, and have ranged from 176 bull trout in 2006 to 1079 bull trout in 2005. We collected tissue and scale samples from all bull trout we handled, and marked all fish with PIT tags, which allowed us to obtain capture histories across years for many fish. We recaptured 53 bull trout that were previously marked ranging between 285 to 1469 days prior. The recaptured bull trout grew an average of 101.4 mm (0.2 mm/day), and gained an average of 1869 g (3.1 g/day). Juvenile bull trout were collected from 15 tributaries within the Kootenai River Basin in British Columbia and Montana to develop a genetic methodology using microsatellite loci to assess whether fish originated above or below Libby Dam. Results indicated that there is a high degree of genetic variation among different bull trout populations within the Kootenai River basin. Jackknife analysis of our baseline data

set indicated that we had a high degree of power (> 95%) to correctly assign unknown fish captured in the Kootenai River as originating either upstream or downstream of Libby Dam. We applied this methodology to the tissue and scale samples collected each year below Libby Dam during the population estimates to predict origin, and estimated that the proportion of the fish originating above Libby Dam ranged from 49.1 percent in 2004 to 62.7 percent in 2006. The majority of the adults assigned to populations above the dam were assigned to the Wigwam River, British Columbia, which represented the tributary with the highest number of bull trout redds in recent years.

PATTERNS OF TROUT SURVIVAL AND MOVEMENT BEFORE AND AFTER LOGGING ON INDUSTRIAL FOREST LANDS

Robert E. Gresswell, U.S. Geological Survey, Northern Rocky Mountain Science Center, Montana State University, Bozeman, Montana 59717

Aaron M. Berger, Department of Fisheries and Wildlife, Oregon State University, Corvallis, Oregon 97331

Douglas S. Bateman, Department of Forest Science, Oregon State University, Corvallis, Oregon 97331

David Hockman-Wert, U.S. Geological Survey – Forest and Rangeland Ecosystem Science Center, 3200 SW Jefferson Way, Corvallis, Oregon 97331

Clear-cut timber harvest continues to be a common practice that frequently results in a patchwork of disturbance across the landscape. Although harvest techniques have greatly improved over the past half-century, effects of contemporary harvest methods on adjacent (point) and downstream (cumulative) portions of the aquatic network are not well documented. Therefore, we sought to quantify spatial and temporal patterns of survival and movement of coastal cutthroat trout (*Oncorhynchus clarkii clarkii*) in two, experimentally-paired watersheds in the Cascade Mountains of Oregon, before and recently after logging. All harvest units were located along non-fish bearing channels upstream from the end of fish distribution. A total of 4406 trout (>100 mm, fork length) were implanted with half-duplex passive integrated transponder (PIT) tags and monitored seasonally during a 5-yr period (3 yrs before and 2 after harvest) using a combination of electrofishing and mobile and stationary PIT-tag antennas. Apparent survival varied widely among seasons and years, and variation among subcatchments was small in comparison. Seasonal apparent survival, regardless of year or subcatchment, was always lowest during the fall (Sep 15-Dec 15). There was no significant effect of this logging treatment on survival of coastal cutthroat trout in this headwater stream network, and although there was some increase in the probability for movement following logging, it remained low. In general, these data suggest that contemporary forest harvest practices regulations provided adequate short-term protection for coastal cutthroat trout from potentially negative consequences of timber harvest in the non-fish bearing portions of the Hinkle Creek drainage.

AVIAN PISCIVORES VECTOR IN THE GREATER YELLOWSTONE ECOSYSTEM

Todd M. Koel, Center for Resources, Fisheries and Aquatic Sciences Program, P.O. Box 168, Yellowstone National Park, Wyoming 82190

Billie L. Kerans, Department of Ecology, Montana State University, P.O. Box 173460, Bozeman, Montana 59717

Scott C. Barras and Katie C. Hanson, USDA/APHIS/WS, National Wildlife Research Center, Mississippi Field Station, P.O. Box 6099, Mississippi State, Mississippi 39762

John S. Wood, Pisces Molecular LLC, 2200 Central Avenue, Suite F, Boulder, Colorado 80301

Although often blamed on movement of trout, the dispersal vector of *Myxobolus cerebralis* among aquatic habitats often remains unknown. Occurrence of whirling disease in native Yellowstone cutthroat trout (*Oncorhynchus clarkii bouvieri*) within the highly protected environment of Yellowstone Lake is one example. Given their local abundances, we sought to clarify the potential role of highly mobile piscivorous birds in the dissemination of *M. cerebralis* to otherwise isolated habitats. Six each of American White Pelicans (*Pelicanus erythrorhynchos*), Double-crested Cormorants (*Phalacrocorax auritus*), and Great Blue Herons (*Ardea herodias*) were held in an aviary and fed known-infected or uninfected rainbow trout (*O. c. mykiss*). Fecal material produced during 10-day periods before and after feeding was collected to determine if *M. cerebralis* could be detected and, if so, remained viable after passage through the gastrointestinal tract of these birds. Fecal samples from all (100%) of the nine birds fed known-infected trout and collected during days 1-4 following feeding tested positive for the presence of *M. cerebralis* by PCR. In addition, *Tubifex tubifex* fed fecal material from known-infected herons produced triactinomyxons in laboratory cultures, confirming the persistent viability of the parasite. Given the infection prevalence of cutthroat trout within Yellowstone Lake, pelicans, cormorants, and herons can move an estimated 1.27 billion *M. cerebralis* myxospores in the ecosystem during a 100-day breeding season each year. Piscivorous birds have the potential to concentrate and release *M. cerebralis* myxospores with fecal material into habitats highly suitable for *T. tubifex*, forming the basis of a positive feedback loop where the proliferation of *M. cerebralis* is supported.

LANDSCAPE CONSERVATION AND THE BIG HOLE CCAA PROGRAM

Nathan Korb, Southwest Montana Director for Science and Stewardship, The Nature Conservancy, 32 S. Ewing Street, Helena, Montana 59601

Beyond the tangible results across riparian and aquatic habitats of the upper Big Hole basin, the CCAA restoration program creates new opportunities for an array of landscape-scale conservation activities. The public and private partnerships forged through the CCAA program and the local watershed groups provide a valuable system for advancing collaboration conservation including: private land conservation, invasive species management, wetland restoration, bird and plant monitoring, fire management, and preservation of other species of concern or federally listed species. Proposed and on-going projects in the Big Hole, as well as lessons that can be applied to other biologically important landscapes, will be presented.

REDUCED GENETIC VARIATION IN UPPER MISSOURI RIVER DRAINAGE WESTSLOPE CUTTHROAT TROUT POPULATIONS APPEARS TO BE DUE TO HISTORICAL AND CONTEMPORARY FACTORS

Robb F. Leary, Montana Fish, Wildlife and Parks, Division of Biological Sciences, University of Montana, Missoula, Montana 59812

Sally Painter, Steve Amish, Angela Lodmell, and Fred W. Allendorf, Division of Biological Sciences, University of Montana, Missoula, Montana 59812

John H. Powell, Hopkins Marine Station, Stanford University, 120 Ocean View Boulevard Pacific Grove, California

The number of westslope cutthroat trout (*Oncorhynchus clarkii lewisi*) local populations has decreased over the past 100 years. In Montana, this decrease has been most severe in the upper Missouri River drainage. The remaining populations in the drainage tend to be small and confined to headwater streams isolated above man made or natural barriers. Data from 14 nuclear loci indicate that mean average expected heterozygosity (Columbia = 0.155, Missouri = 0.064), mean proportion of polymorphic loci (0.321, 0.179), and mean average number of alleles per locus (2.040, 1.313) were all significantly smaller within 16 Missouri populations than within 34 upper Columbia River populations. Total heterozygosity among the Missouri populations (0.106) was only about half of that observed among 16 randomly chosen Columbia populations (0.194). The average number of alleles per locus was also lower among the Missouri (2.714) than Columbia (3.643) populations. These latter two observations suggest that the reduced genetic variation in the Missouri populations is partially the result of a significant founder effect when the fish colonized the drainage. The relative amount of genetic divergence among the Missouri populations ($F_{ST} = 39.4\%$) was about twice that observed among the Columbia populations (20.5%) suggesting that subsequent to colonization the former have experienced more genetic drift and isolation than the latter. Because of their reduced genetic variation, genetic rescue is more likely to be required in Missouri than Columbia populations. Furthermore, the reduced genetic variation in the Missouri populations may retard their response to other conservation actions.

A FISH HATCHERY'S ROLE IN A CHANGING MONTANA

Mark G. Maskill, USDI Fish and Wildlife Service, Creston National Fish Hatchery, Kalispell, Montana 59901

Demographic change is sweeping across Montana like a racing wild fire, rapidly changing the mix of rural and urban dwellers and the face of Montana politics. A good percentage of these new residents have relocated to Montana specifically to partake in the time honored Montana traditions of hunting and fishing. Interest in fish and wildlife is at an all time high. As the natural resources of our state are pursued more intensely by residents and nonresidents, as well as consumptive and non-consumptive users, the management of these resources must adjust in order to protect fragile native populations. How will fish managers respond to this growth? What role should fish hatcheries play in Montana today? Are fish managers utilizing this conservation tool in the right manner? How much effort should be put into restoring threatened native populations? What attempts should be made trying to increase recreational opportunities throughout the state? The Creston NFH has served a unique role in Montana fish production, constantly changing and evolving, for over seventy years. Creston

has played a support role in several restoration and recreational fishery projects. Some of these have failed and some have created world class fisheries. This talk will focus on the lessons we have learned and how we can put those lessons to use in playing a future role in conserving, protecting, and enhancing Montana's fishery resource.

HYBRIDIZATION RAPIDLY REDUCES FITNESS OF NATIVE CUTTHROAT TROUT IN THE WILD

Clint C. Muhlfeld*, U.S. Geological Survey, Northern Rocky Mountain Science Center, Glacier Field Office, Glacier National Park, West Glacier, Montana 59936

Steven T. Kalinowski, Thomas E. McMahon, and Mark L. Taper, Montana State University, Department of Ecology, Bozeman, Montana 59717

Sally Painter and Fred W. Allendorf, University of Montana, Conservation Genetics Laboratory Missoula, Montana 59812

Robb F. Leary, Montana Fish, Wildlife and Parks, University of Montana, Missoula, Montana 59812

Human-mediated hybridization between native and invasive species is a leading cause of biodiversity loss worldwide. How hybridization affects fitness and what level of hybridization is permissible pose difficult conservation questions with little supportive empirical information to inform policy and management. This is particularly true for salmonids, where widespread introgression among nonnative and native taxa often create hybrid swarms over extensive geographic areas, threatening natives with genomic extinction. Here, we used parentage analysis with multilocus microsatellite markers to measure how varying levels of genetic introgression with nonnative rainbow trout (*Oncorhynchus mykiss*) affect reproductive success (number of offspring per adult) of native westslope cutthroat trout (*O. clarkii lewisi*) in the wild. Small amounts of hybridization markedly reduced reproductive success of male and female trout, with fitness sharply declining by ~50 percent with only 20 percent nonnative admixture. Despite heavy fitness costs, our data suggest that hybridization may spread due to relatively high fitness of first-generation hybrids and inordinately high reproductive success of a few males with high levels of nonnative admixture. This outbreeding depression suggests that even low levels of nonnative genetic admixture may have negative effects on reproductive success in the wild and that policies protecting hybridized populations may need reconsideration.

EFFECTS OF LARGE WOOD PLACEMENT ON CHANNEL MORPHOLOGY AND AQUATIC HABITAT HALLOWAT CREEK, NORTH FORK FLATHEAD RIVER DRAINAGE, MONTANA

John Muhlfeld* and Jonathan Ferree, River Design Group, Inc., 5098 Highway 93 South, Whitefish, Montana 59937

Montana Fish, Wildlife and Parks (MFWP) documented a substantial reduction in bull trout (*Salvelinus confluentus*), spawning redds in Hallowat Creek, a tributary to Big Creek of the North Fork Flathead River. The Big Creek watershed is an important spawning tributary for fluvial bull trout, a federally-listed threatened species (USDI Fish and Wildlife Service 1998). USDA Forest Service fire suppression efforts undertaken during the 2001 Moose Creek

Fire influenced channel hydraulics and spawning habitat distribution in Hallowat Creek. Fire suppression activities included cutting large woody debris within the active channel and riparian zone. Stable large wood loss resulted in a more simplified channel characterized by coarse substrate, few pools, and infrequent large woody debris. In 2005, River Design Group, Inc. (RDG), MFWP and FNF implemented an in-channel treatment plan for Hallowat Creek. Large wood was imported to the active channel and arranged in stable wood complexes to promote pool development. Reach-specific treatments were developed to either meet or exceed large woody debris counts that were enumerated in a pre-fire, 1998 R1 R4 survey conducted by the FNF. Channel monitoring surveys and annual bull trout redd counts were completed in 2005, 2006 and 2008 to assess spawning and geomorphic response to the structures. Results suggest augmenting existing stable large wood structures with additional wood, and building structures that are anchored to stable floodplain features provide the best opportunity for increasing complex pool habitat in Hallowat Creek. This presentation summarizes the monitoring data and provides recommendations to practitioners engaged in similar habitat restoration projects in forested, mountain streams.

CREATING WILDLIFE HABITAT AND EXPANDING FISHERIES WITH BIOHAVEN FLOATING TREATMENT WETLANDS

Tim Mulholland P.E. and Christine Pierce, Headwaters Floating Island, 1018 2nd Ave. North Billings, Montana 59101

BioHaven[®] floating islands are an environmentally low impact and economical way to create new wetland, while augmenting the vitality and biodiversity of a waterway. They are made using non-toxic recycled materials and are hands off and non-mechanical. The biomimetic design gives our fish, wildlife and waters the least carbon footprint and the most natural support for the systems they depend upon for healthy and productive lifecycles. The benefits are many: habitat creation and restoration for a myriad of fauna, from microbiological systems to aquatic and wetland wildlife; reduced turbidity and pollution mitigation (with resource recovery of nutrients such as nitrate and phosphate); and aesthetic and unique living water features of any size or shape for any water body. Floating Islands used in landscaped water features and pools offer a chemical free approach to managing water quality. Placed in a stream or pond, BioHavens and their suspended root complex give secure habitat for fish, offering shade and protection from currents and predators. The living islands attract a range of insects that also become a food source for fish and wildlife and further promote biodiversity. Unique configurations can be designed for nesting sites, spawning platforms and load bearing hanging banks to increase protective fish habitat. Several projects have developed successful, protective bird habitat specifically for Duck, Loon and Swan and a 22,000 sf Caspian Tern Habitat Island.

AN OVERVIEW OF THE CANDIDATE CONSERVATION AGREEMENT WITH ASSURANCES PROGRAM, WITH SPECIAL REFERENCE TO THE BIG HOLE RIVER, MONTANA

Douglas P. Peterson*, USDI Fish and Wildlife Service, Division of Ecological Services, 585 Shepard Way, Helena, Montana 59601

In 1999 the Candidate Conservation Agreement with Assurances (CCAA) program was initiated by the USDI Fish and Wildlife Service to engage private landowners in species conservation before a potential listing under the Endangered Species Act (ESA). For their

participation, private landowners receive assurances that they will not be subjected to additional regulatory burden should the species be listed. Hundreds or thousands of species are potentially eligible for this program, but only a few dozen CCAA's have been completed and the scope and complexity varies widely among individual agreements. Because the program is relatively new, there are little data to evaluate the biological effectiveness of CCAAs. However a few agreements can be considered a success in terms of landowner participation. Issuing permits under the ESA has led to litigation for other conservation programs like Habitat Conservation Plans (HCPs), but the validity of the CCAA policy and individual CCAA agreements have not yet been subjected to formal legal challenge. I briefly summarize the basic premise of the CCAA program, the different organizational frameworks to structure these agreements, and some of the institutional and social challenges to effective implementation, highlighting the example of the CCAA for fluvial Arctic grayling in the upper Big Hole River, Montana.

GENETIC VARIATION, ANCESTRY AND POPULATION STRUCTURE IN NATIVE ARCTIC GRAYLING IN THE UPPER MISSOURI RIVER

Douglas P. Peterson, USDI Fish and Wildlife Service, Division of Ecological Services, 585 Shepard Way, Helena, Montana 59601

William R. Ardren, USDI Fish and Wildlife Service, Lake Champlain Fish and Wildlife Resources Office, 11 Lincoln Street, Essex Junction, Vermont 05452

Arctic grayling (*Thymallus arcticus*) were thought to be historically widespread in the Missouri River system in Montana and Wyoming, but have been reduced to a handful of remnant populations since Euro-American colonization. Conservation efforts have focused primarily on protecting the fluvial population in the Big Hole River, and re-establishing fluvial populations elsewhere. Widespread historical stocking of exogenous grayling has created some uncertainty about the origin of extant populations and the composition of native gene pools. Additionally, declines in native populations may have reduced the genetic template for adaptation and recovery. Effective conservation would benefit from a better understanding of genetic ancestries, and whether bottlenecks have substantially altered genetic diversity of remnant populations. Consequently, we conducted a population-level genetic analysis of native and introduced grayling from 18 locations. We genotyped 730 grayling at 10 microsatellite loci and used these data to identify population groupings, genetic variation within and among groups, and evaluate evidence for population bottlenecks. We found significant divergence among native populations from the Big Hole River, Madison River, and Red Rock lakes. The Big Hole population had greater heterozygosity and allelic diversity than the Madison and Red Rock populations, both of which showed some evidence of recent bottlenecks. Most introduced populations traced their ancestry to the adfluvial Red Rock lakes population, and we did not find strong evidence that stocking of hatchery grayling homogenized native gene pools. Geographic patterns of genetic variation among native Missouri River grayling were consistent with differentiated local populations historically connected by occasional gene flow.

THE EAST FORK SPECIMEN CREEK WESTSLOPE CUTTHROAT TROUT RESTORATION PROJECT: THE GOOD, THE BAD, AND THE SMOKEY

Michael E. Ruhl, Todd M. Koel, Jeffery L. Arnold, and Brian D. Ertel, Yellowstone Center for Resources, Fisheries and Aquatic Sciences Section, P.O. Box 168, Yellowstone National Park, WY 82190

Initiation of the East Fork Specimen Creek (EFSC) westslope cutthroat trout (*Oncorhynchus clarkii lewisi*; WCT) restoration project in 2005 marked the beginning of a renewed effort to actively restore native fish in Yellowstone National Park. That summer an interdisciplinary team was assembled to begin a NEPA compliance process and within 12 months an EA and FONSI were complete. Project implementation began immediately with chemical removal of nonnative fish in High Lake, a 7-ac isolated headwater lake, in August 2006. Unfortunately, the construction of a fish barrier on EFSC, planned for completion in 2007, was delayed by a natural wildfire that destroyed all previous work and left an unsafe worksite. Rotenone treatment of High Lake was successful and we began efforts to restock the lake with genetically-pure WCT (from multiple sources) that year. In 2008 we again stocked High Lake with WCT and undertook a vigorous effort to complete the EFSC fish barrier. Because of its remoteness, 93 mule loads and five helicopter sling loads were required to move supplies and tools to the site. Contracted log crafting specialists, Montana Conservation Corps crews, and park staff collaborated to bring the barrier to successful completion. Immediately afterwards two piscicide treatments were conducted on EFSC from High Lake downstream to the fish barrier (12 km). Additional treatments of this reach are planned for 2009 and WCT reintroduction efforts will begin as soon as complete removal of non-native fish is verified.

CLIMATE CHANGE MEDIATES THE SPATIAL PARTITIONING OF SCULPIN AND LONGNOSE DACE LEADING TO TROPHIC CASCADES IN RIVERINE ECOSYSTEMS OF WESTERN MONTANA

David A. Schmetterling and Robert Clark, Montana Fish, Wildlife and Parks, 3201 Spurgin Road, Missoula, Montana 59804

Susan B. Adams, USDA Forest Service, Southern Research Station, 1000 Front Street, Oxford, Mississippi 38655

Michael K. Young, Rocky Mountain Research Station, 800 East Beckwith Avenue, Missoula, Montana 59801

Sculpin (*Cottus spp.*) and longnose dace (*Rhinichthys cataractae*), perform diverse roles in trophic interactions. Both are abundant inhabitants of the benthos, and longnose dace are present in nearly every Montana drainage. The ecological importance of both genera arises, in part, from their tendency to occur in high abundances, often dominating fish assemblages in number and biomass. Although both are numerous, small-bodied, and largely confined to the benthos, they occupy different trophic positions, express different life history tactics, and perform different ecological roles. Over the last few decades, west-central Montana rivers have warmed affecting the distribution of, among other species, longnose dace, and sculpin. Both species have undergone dramatic shifts in occupancy of habitats leading to ecotonal shifts on a broad geographical scale and this change has occurred rapidly and quietly. In general, as waters have warmed, longnose dace have replaced sculpin leading to the retreat and isolation of many sculpin metapopulations. Both species are sympatric with endangered species like the bull trout (*Salvelinus confluentus*), whose populations are targets for

protection, mitigation, and enhancement. It is unclear the effects of the replacement of sculpin by longnose dace, though they are likely large and measurable, and since they represent the basis of the food chain for higher trophic order organisms, this change could ultimately could affect the restorative potential of the systems for target species. Furthermore, warming and species replacement may lead to a further homogenization or regional distinctive ecotones and fauna.

WATERSHED-SCALE APPROACH TO ASSESSING COLORADO RIVER CUTTHROAT TROUT ABUNDANCE AND HABITAT IN THE UPPER COLORADO RIVER HEADWATERS

Ted R. Sedell, Montana State University, PO Box 173460, Bozeman, Montana 59717

Robert E. Gresswell, U.S. Geological Survey, NoROCK, Montana State University, 229 Johnson Hall, Bozeman, Montana 59717

Information concerning the effects of physical habitat on the distribution, abundance, and population structure of salmonid fishes has been routinely gathered at the site and reach scales, but data collected at these scales has provided little insight into spatial and temporal structure of fish assemblages within the stream network. This is especially relevant where periodic disturbances across the landscape are extensive, and effects are not uniform (spatially or temporally). To gain new insight into the relationships between trout abundance and population structure and the physical habitat at multiple spatial scales, we surveyed six watersheds in the upper Colorado River basin and assessed Colorado River cutthroat trout (*Oncorhynchus clarki pleuriticus*) abundance and habitat variables continuously throughout the stream network. The watersheds were selected randomly from a sampling frame based on presence of Colorado River cutthroat trout populations, percent beetle infestation of the watershed area, and erosion potential of the upslope geology. Although similar physical habitat conditions existed within and among several watersheds, fish distribution was patchy at the channel-unit scale, and physical habitat was not strongly related to distribution. However, at the reach scale, patterns of fish distribution explained a substantial amount of the variation in abundance. Further refining of these relationships will provide critical information necessary to assess future changes related to restoration activities, wildfire, and climate change.

RECOVERY OF WESTSLOPE CUTTHROAT TROUT POPULATIONS FOLLOWING REMOVAL OF NONNATIVE BROOK TROUT

Bradley B. Shepard, Montana Cooperative Fishery Research Unit, Ecology Department, Montana State University, Bozeman, Montana 59717-3460 and Montana Department of Fish, Wildlife and Parks, 1420 East Sixth Avenue, Helena, Montana 59620-0701

Mark L. Taper, Ecology Department, Montana State University, Bozeman, Montana 59717

Alexander V. Zale, Montana Cooperative Fishery Research Unit, U.S. Geological Survey, Montana State University, Bozeman, Montana 59717-3460

We investigated whether 75-mm and longer westslope cutthroat trout (*Oncorhynchus clarkii lewisi*) and brook trout (*Salvelinus fontinalis*) occupied similar niches by comparing biomasses, population densities, and condition factors prior to and following total removal of brook trout in 2.3- to 3.0-km reaches of three headwater streams in Montana. Biomasses and their associated errors were estimated using depletion estimators. Total trout biomass did not

change significantly after brook trout removal indicating that these two species have similar niches in these streams. Densities of juvenile westslope cutthroat trout were significantly and negatively affected by densities of juvenile brook trout and positively related to densities of adult westslope cutthroat trout ($R^2 = 0.482$; F-ratio = 15.415; $P < 0.001$). Including densities of westslope cutthroat trout or brook trout from the previous year did not measurably improved model performance. Densities of juvenile brook trout were negatively associated with body condition of juvenile westslope cutthroat trout. We found evidence for size-asymmetric competition in one stream, but not in the other where it was assessed. Interspecific competition between brook trout and westslope cutthroat trout was nearly as strong as intraspecific competition within westslope cutthroat trout, especially among juveniles, providing insight into one mechanism by which brook trout displace westslope cutthroat trout.

SPATIAL AND TEMPORAL FISH ENTRAINMENT FROM HAUSER RESERVOIR, MONTANA

Justin P. Spinelli, Montana Cooperative Fishery Research Unit, Department of Ecology, Montana State University, P.O. Box 173460, Bozeman, Montana 59717-3460

Alexander V. Zale, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Department of Ecology, Montana State University, P.O. Box 173460, Bozeman, Montana 59717

Management sport fish populations of Hauser Reservoir, Montana, is hindered by undesirable and unpredictable downstream entrainment of fish through Hauser Dam. We quantified fish entrainment through Hauser Dam using hydroacoustic technology at turbine intakes from July 2007 to November 2008 and over the spillway from 21 May to 18 July 2008. Species composition of entrainment was characterized using multiple netting gears. Annual estimated turbine entrainment was higher in 2007 ($N = 99,148 \pm 5582$) than in 2008 ($N = 53,456 \pm 5,118$). Spillway entrainment ($N = 29,931 \pm 3173$) was 36 percent of total annual entrainment in 2008. Entrainment was higher in fall than in summer in both years, likely in response to fall turnover and the annual release of hatchery rainbow trout. Most entrained fish (~60%) were < 220 mm total length in both turbine discharge and spill. The most common fish captured were rainbow trout (43%), white sucker (27%), and walleye (15%). The least common were common carp and yellow perch. We applied species composition by size to the hydroacoustic data to identify fish species entrained, but most fish ($N = 74,062 \pm 4834$) could not be reliably assigned to a species because concurrent net catches did not include individuals of similar size. Most identified entrained fish were rainbow trout ($N = 33,472 \pm 3014$) and walleye ($N = 35,439 \pm 2953$). Identification of patterns in spatial and temporal fish losses affords fishery managers the ability to make more informed decisions about operation of this dam.

MERCURY CONTAMINATION IN THE FISHES OF GLACIER NATIONAL PARK

Craig Stafford, P.O. Box 1472, Missoula, MT 59806

Chris Downs, Science Center - Glacier National Park, West Glacier, Montana 59936

Heiko Langner, Department of Geosciences - University of Montana, 32 Campus Drive Missoula, Montana 59812

Elizabeth McGarry, Department of Biology - St. Thomas University, 2115 Summit Ave. St. Paul, Minnesota 55105

We investigated mercury contamination in fishes from four lakes in Glacier National Park. Three of these lakes (Bowman, Harrison, McDonald) are on the west side of the park, and one is on the east side (St Marys). We focused our sampling on lake trout (*Salvelinus namaycush*) but also collected lake whitefish (*Coregonus clupeaformis*), burbot (*Lota lota*), and incidentally killed bull trout (*Salvelinus confluentus*). Mercury contamination generally increased with fish size but not always. Lake trout and burbot were the most contaminated species and levels exceeding 0.5 ppm Hg wet occurred in the larger fish. We found that lake whitefish and bull trout generally had lower mercury levels than lake trout on a size normalized basis. Examination of the lake trout and lake whitefish data revealed that males and females had similar levels of mercury. Mercury levels in lake trout from the west side lakes showed similar trends with fish size, and were comparable to other lake trout populations in the region.

ADJUSTING LAKE TROUT AGES VERSUS OTOLITH MASS RELATIONSHIPS FOR VARIABLE GROWTH

Craig Stafford, Division of Biological Sciences - University of Montana, 32 Campus Drive Missoula, MT 59812

Dale Hanson, Green Bay National Fish & Wildlife Conservation Office - USDI Fish and Wildlife Service, 2661 Scott Tower Drive, New Franken, Wisconsin 54229

Otolith weight holds considerable promise for increasing speed and consistency of age estimates relative to otolith thin sections, especially for species such as lake trout (*Salvelinus namaycush*) where age and otolith weight are highly related. Using known age lake trout from Lake Michigan we demonstrate that the age versus otolith mass relationship is dependent body growth rate, reducing the utility of otolith mass alone for estimating age when growth rates vary between samples. To compensate for this growth bias we developed a correction procedure. This procedure uses otolith weight versus total length relationships to estimate the body growth rate difference between samples, and then estimates the bias in the age versus otolith mass relationship. Using the correction procedure we successfully adjusted the Lake Michigan age versus otolith mass relationship to the more slowly growing Flathead Lake fish, revealing a pattern of declining body growth rates through time in the Flathead Lake population.

ARE LAKE TROUT IN FLATHEAD LAKE MORPHOLOGICALLY AND GENETICALLY SEGREGATED BY DEPTH?

Craig Stafford, Lisa Eby, and Fred Allendorf, University of Montana, 32 Campus Drive Missoula, Montana 59812

Megan McPhee, Flathead Lake Biological Station, University of Montana, 32125 Bio Station Lane, Polson, Montana 59860

We compared muscle lipid content, muscle stable isotope ratios, body morphology, and microsatellite allele frequencies between shallow (0-25 m) and deep caught (>50 m) lake trout (*Salvelinus namaycush*) from Flathead Lake, Montana. We found that lipid content was similar between depth groups. Stable isotopes of N and C varied between depth groups, demonstrating that individual fish exhibit long term depth preferences. Depth groups varied in their morphology. Relative to shallow fish, deep fish had a head more in line with the rest of the body, bigger eye, deeper body, wider skull, longer pectoral fin, and a deeper and shorter caudal peduncle. Microsatellite allele frequencies were similar between depth groups, strongly suggesting gene flow between shallow and deep fish.

WESTERN PEARLSHELL MUSSEL DISTRIBUTION & STATUS IN MONTANA: TWO YEARS LATER IT'S WORSE THAN WE THOUGHT!

David M. Stagliano, Montana Natural Heritage Program, 1515 East 6th Avenue, Helena, MT 59620

Montana's only trout stream mussel, the western pearlshell (*Margaritifera falcata*), has disappeared from many of our watersheds in relatively recent times. During 2006 and 2007, we reviewed western pearlshell occurrence records and systematically resurveyed sites of current and historic occupation in watersheds throughout the state. Twenty-five of the original 40 site records proved to be either absent or had non-viable mussel populations; only seven of the 15 viable populations documented in 2007 were classified as having excellent integrity. Extensive field surveys and biologist training workshops continued in 2008, but only four additional viable populations were reported. Of the ~820 stream reaches (avg. length ~150 m) surveyed over the course of 3 yrs, western pearlshell populations were absent from 660 (76 %) of the reaches and non-viable to severely declining from another 139 sites (16%). Sites with excellent population viability (9) were rare and represented disjunct **metapopulations** with little ability to colonize other stream reaches in the watershed. **Because of this fact**, the evaluation of current vs. previously occupied river miles and the severity of the decline, we officially placed this species on the Species of Concern list in November ranked an S2 (vulnerable to extirpation in the state). Introduction of **non-native** fishes, reduction of westslope cutthroat populations, reduced in-stream flows and **warmer water** temperatures have all been implicated in the decline of populations of the western **pearlshell**. Reintroduction of western pearlshells into westslope cutthroat trout restored stream reaches is currently being investigated.

PROPOSED COAL MINING AND COAL-BED METHANE DEVELOPMENT THREATEN AQUATIC RESOURCES IN THE TRANSBOUNDARY FLATHEAD ECOSYSTEM

Amber C. Steed, Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, Montana 59901

Clint C. Muhlfield, U.S. Geological Survey, Northern Rocky Mountain Science Center, Glacier Field Office, Glacier National Park, West Glacier, Montana 59936

Erin K. Sexton, Research Scientist, Flathead Lake Biological Station, University of Montana, 32125 Bio Station Lane, Polson, Montana 59860-6815

The Transboundary Flathead River basin in Montana (USA) and British Columbia (Canada) hosts one of the most diverse and unique aquatic ecosystems throughout North America. Migratory bull trout (*Salvelinus confluentus*) and non-hybridized westslope cutthroat trout (*Oncorhynchus clarkii lewisi*) migrate from Flathead Lake upstream to the Canadian headwaters to spawn and rear, representing some of the last remaining strongholds in the basin. However, proposed open-pit coal mining and coalbed methane (CBM) drilling in the Canadian headwaters threaten water quality, invertebrate communities, and migratory fish populations downstream to Glacier National Park (GNP) and Flathead Lake. In response to these threats, a multi-agency, long-term research and monitoring program was initiated in 2005 to examine water and sediment chemistry, contaminant levels, aquatic habitat, and the distribution and genetic diversity of native fishes. Comparative data collected in the neighboring Elk River drainage, a system impacted by coal mining and CBM development, show increased nutrients and heavy metal concentrations in the water and lower invertebrate species richness than the Flathead. In 2008, basin-wide fisheries surveys were initiated and data were collected at 119 sites in Canada and GNP. Native fishes were found throughout much of the system, including proposed mining locations. Additionally, the highest quantity of bull trout redds in the system was detected immediately downstream of proposed mine sites. Continuation of these collaborative investigations will provide necessary baseline data to inform conservation and management decisions impacting this diverse and sensitive transboundary system.

ANALYSIS OF POPULATION METRICS TO ASSESS THE EFFICACY OF LAKE TROUT SUPPRESSION IN YELLOWSTONE LAKE, YELLOWSTONE NATIONAL PARK

John M. Syslo and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Lewis Hall, Montana State University, Bozeman, Montana 59717

Patricia E. Bigelow, Philip D. Doepke, and Todd M. Koel, Center for Resources, Fisheries and Aquatic Sciences Program, P.O. Box 168, Yellowstone National Park, Wyoming 82190

Introduced lake trout (*Salvelinus namaycush*) threaten to extirpate native Yellowstone cutthroat trout *Oncorhynchus clarkii bouvieri* from Yellowstone Lake, Yellowstone National Park. The USDI National Park Service removed nearly 280,000 lake trout from Yellowstone Lake between 1997 and 2007. Lake trout population size has not been estimated; therefore, it is difficult to determine what proportion has been removed. We evaluated several population metrics to determine if the removal program caused the lake trout population to exhibit characteristics typical of overharvested populations. Biomass of lake trout harvested has

increased through the duration of the suppression program and was 0.74 kg/ha in 2007. Catch-per-unit-effort of lake trout has increased in both targeted-removal and survey netting. Mean length at age declined from 2000 through 2006 for lake trout older than 8 years. Total annual mortality in 2007 was 37 percent for lake trout of ages 3 through 5 and 12 percent for lake trout over age-5. Population metrics do not indicate that lake trout suppression has been effective at reducing abundance in Yellowstone Lake, however, the quality of existing data limit the certainty of conclusions. Yield-per-recruit modeling indicates that growth overfishing may begin to occur when conditional fishing mortality exceeds 10 percent for lake trout of all ages. Spawning potential ratio computed with rates of mortality from 2007 was 0.14, and could be reduced to 0.04 with a conditional fishing mortality rate of 20 percent for lake trout over age-5. An increase in fishing mortality of lake trout over age-5 may increase effectiveness of lake trout suppression.

DETERMINING MORPHOLOGICAL AND BIOCHEMICAL PARAMETERS ASSOCIATED WITH EARLY OVARIAN FOLLICULAR ATRESIA IN WHITE STURGEON FEMALES

Mariah J. Talbott and Christopher S. Guy, Montana Cooperative Fishery Research Unit, Montana State University, Bozeman, Montana 59717

Joel P. Van Eenennaam, Javier Linares-Casenave, and Serge I. Doroshov, Department of Animal Science, University of California, Davis, California 95616

Molly A.H. Webb, Bozeman Fish Technology Center, 4050 Bridger Canyon Road, Bozeman, Montana 59715

In order to improve quality and yield of caviar in farmed white sturgeon (*Acipenser transmontanus*), it is essential to correctly assess stage of ovarian maturity and avoid harvesting females with atretic ovarian follicles. To detect atresia by changes in blood plasma parameters, individual females ($N = 11$) in the late phase of oogenesis were repeatedly bled and their ovaries biopsied before and after onset of ovarian atresia. Follicular atresia was induced by transferring females at Sterling Caviar, LLC, California from cold (10–13 °C) to warm water (20 °C). Follicle diameter increased and oocyte polarization indices decreased over time. Plasma testosterone and estradiol concentrations in fish with normal follicles were higher, compared to fish exhibiting early histological signs of follicular atresia, such as structural changes in the egg coat. Total plasma protein and calcium concentrations did not differ between fish with normal and regressing ovaries. In the future, our study may benefit sturgeon conservation propagation programs in Montana by improving techniques for detection of ovarian atresia in the late phase of oogenesis.

DISTRIBUTION, ABUNDANCE, AND AGE STRUCTURE OF JUVENILE BULL TROUT IN A TRIBUTARY TO QUARTZ LAKE, GLACIER NATIONAL PARK, MONTANA

Lora B. Tennant and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Montana State University, PO Box 173640, Bozeman, Montana 59717-3460

Robert E. Gresswell, U.S. Geological Survey, Northern Rocky Mountain Science Center, Montana State University, P.O. Box 173492, Bozeman, Montana 59717

Lacustrine-adfluvial bull trout (*Salvelinus confluentus*) occupy lakes west of the Continental Divide in Glacier National Park (GNP), Montana. Research in GNP has focused primarily on the relative abundance of bull trout within lakes, patterns of connectivity among bull trout populations, and interactions between adult bull trout and nonnative lake trout (*S. namaycush*). Consequently, there is little known about the ecology, abundance, and distribution of juvenile bull trout within headwater drainages in GNP. The expanding distribution and potential negative effects of lake trout on bull trout within GNP has made understanding the ecology of juvenile bull trout a high priority. This study documented the distribution, relative abundance, and age structure of juvenile bull trout in the Quartz Drainage upstream of Quartz Lake. The study area included Quartz Creek and Rainbow Creek; a tributary that enters Quartz Creek about 1.25 km upstream of Quartz Lake. Juvenile bull trout were sampled using a backpack electrofishing unit; bull trout were enumerated and measured for length. Juvenile bull trout were distributed throughout the study area. Relative abundance of juvenile bull trout was greater downstream of the confluence of Quartz and Rainbow creeks than upstream. A bimodal length-frequency distribution suggested that the bull trout present were age-1 and age-2+. These data provide baseline information needed for future recovery efforts aimed at mitigating potential negative effects associated with lake trout colonization.

MONTANA UNAUTHORIZED FISH INTRODUCTIONS

Jim Vashro, Regional Fisheries Manager, FWP, 490 N. Meridian, Kalispell, Montana 59901

Montana Fish, Wildlife and Parks has maintained an unauthorized fish introduction database for more than 15 years that documents unwanted fish introductions across the state that have persisted long enough to be detected. The database currently covers 49 species of fish in 536 introductions in 298 waterbodies. Unauthorized introductions cover every drainage in the state and have caused impacts on native and recreationally important fish species, increased management costs, decreased recreational opportunity, and spread disease. Types and spread of introductions are discussed. The motivations behind illegal introductions are examined and linked to possible means of prevention and control.

LIFETIME MOVEMENT PATTERNS OF CUTTHROAT TROUT IN A STREAM NETWORK

Michael K. Young, Rocky Mountain Research Station, 800 East Beckwith Avenue, Missoula, Montana 59801

Although recent research has emphasized the ubiquity of movements in freshwater fish life histories, the timing, prevalence, and extent of these movements remain contentious. I used juvenile-adult distributions, PIT tagging, and turnover rates of unmarked fish to assess lifetime movement patterns of Colorado River cutthroat trout throughout the 40-km North

Fork Little Snake River watershed in southern Wyoming from 1996 to 2001. Turnover rates indicated that about 30 percent of fish moved an average of at least 100 m annually. Juvenile and adult fish exhibited little spatial overlap in about half of the stream segments in this basin, and length-frequency discrepancies between tributary and main-stem fish implied widespread movement of fish age-2 and younger. About 40 percent of recaptured PIT-tagged fish were mobile. The probability of movement was related to the length of time between first and last capture and to specific growth rate, but not to fish length or condition. Fish marked in tributaries were less likely to move and moved shorter distances than did fish marked in the main stem, but fish from both sources were most likely to move upstream between captures. Median movement was 300 m and many fish moved several kilometers and between tributaries and the main stem, but migration barriers appeared to influence movement in some areas. These complex movement patterns support the conclusion that even slow-growing salmonids in small streams regularly alter their positions to feed, grow, reproduce, and seek refuge from unfavorable environments.

Aquatic Poster Session

RESPONSE OF NON-TARGET ORGANISMS FROM ROTENONE TREATMENTS WITHIN THE EAST FORK SPECIMEN CREEK DRAINAGE, YELLOWSTONE NATIONAL PARK

Jeffrey L. Arnold, Todd M. Koel, Brian D. Ertel, and Michael E. Ruhl, Yellowstone Center for Resources, Fisheries and Aquatic Sciences Section, P.O. Box 168, Yellowstone National Park, Wyoming 82190

Fish removal by rotenone applications is a highly effective fish management tool. However, rotenone is nondiscriminatory and can have negative impacts on non-target aquatic organisms. In 2004 Yellowstone National Park staff began planning a native fish restoration project within East Fork Specimen Creek. We conducted pre- and post- treatment monitoring on invertebrate communities throughout the drainage and on amphibian populations in the vicinity of High Lake, a 7-ac headwater lake where initial rotenone treatments took place. In August 2006 fish in High Lake were chemically removed using rotenone (CFT-Legumine). CFT Legumine is a relatively new formulation of rotenone in the United States that doesn't contain petroleum hydrocarbon solvents that are used in traditional rotenone formulas. As a result, CFT Legumine is likely less harmful to the environment but affects on non-target organisms, such as invertebrates and amphibians, are poorly understood. Among invertebrate populations, both pre- and post- treatment studies indicated that midge larvae were the most common invertebrate groups in the stream and lake benthos with increasing densities after rotenone treatment. Results from the stream invertebrate samples indicate that mayfly, stonefly, and caddisfly larvae were most susceptible to rotenone with some taxa experiencing 100-percent mortality. One year after treatment, however, most taxa had recovered with densities exceeding pre-treatment conditions. Higher invertebrate densities could be a result of the absence of fish predation the year following treatment. Similarly, larval amphibians appeared to experience 100-percent mortality from the initial rotenone application but tadpoles were observed in greater numbers 1 yr post-treatment.

IMPACTS OF TWO SMALL DAMS ON MIGRATORY BULL TROUT IN THE CLEARWATER RIVER, MONTANA

Aubree Benson, and Lisa Eby, University of Montana, Wildlife Biology Program, 32 Campus Drive, Missoula, Montana 59812

Ladd Knotek and John Thabes, Montana Fish, Wildlife and Parks, 3201 Spurgin Road, Missoula, Montana 59804

Dams are well known for their negative impacts on fish populations, and dam removal decisions are becoming increasingly common. In collaboration with Montana Fish, Wildlife and Parks and the USDA Forest Service, we used radio telemetry to explore the impacts of the small Emily-A and Rainy Dams on movement of migratory bull trout (*Salvelinus confluentus*) throughout the Clearwater River Drainage. We captured a total of 88 adfluvial bull trout or bull trout/brook trout (*S. fontinalis*) hybrids below the two small dams, primarily by angling. We implanted radio tags in 31 fish and released them above the dams, passing a total of 75 fish in 2007-2008. We monitored their movements and those of 27 other bull trout tagged in the surrounding lakes. The Emily-A is a complete upstream migration barrier, whereas Rainy is a partial barrier. Ninety-seven percent of the radio tagged fish we moved over the dams swam into one of three previously unknown spawning tributaries and presumably spawned. Although we passed a relatively large number of bull trout, redd counts were low, and we estimated only 13 percent detection probability at the dams. In one tributary, approximately 40-47 percent of the spawning adults were fish we passed over the Emily-A Dam. Post-spawning mortality rates were high and attributed primarily to predation. Our data suggests that the dams have large impacts on population sustainability, and will contribute to the decision-making process involving dam modifications or removal to balance the benefits of upstream passage for native fish with the risk of expansion by undesirable non-native fish.

MANAGEMENT OF ANTHROPOGENICALLY DERIVED HYBRID POPULATIONS: EXPLICIT RECOGNITION OF ASSUMPTIONS

Matthew P. Corsi, Wildlife Biology Program, University of Montana, 32 Campus Drive, Missoula, Montana, 59801

Paul Spruell, Department of Biology, Southern Utah University, 351 W. University Boulevard, Cedar City, Utah 84720

Charles E. Corsi, Idaho Department of Fish and Game, 2885 Kathleen Avenue, Coeur d'Alene, Idaho 83815

Hybridization and introgression between native and introduced species is one of the most challenging issues currently facing fisheries managers. While recognizing we are simplifying arguments, we suggest two hybrid management paradigms have emerged. The first posits that as long as introgression is at moderate to low level, and the resulting hybrids are morphologically and ecologically similar to the native taxon, they should be considered a member of the parental species. The alternative view suggests that conservation efforts should be focused on pure native genomes that have evolved in response to localized selective pressures and hybridized populations are a conservation threat. We suggest that both management approaches are based on a few key assumptions about the nature and ultimate outcome of hybrid fitness and ecology. Although these assumptions are implicit in the arguments presented by both sides of the debate, neither the assumptions nor the management

implications of violations of those assumptions are consistently clarified and discussed. In our poster, we present a framework that addresses various assumptions surrounding hybridization in cutthroat trout (*Oncorhynchus clarkii*) populations by introduced rainbow trout (*O. mykiss*) and the ecological outcomes each assumption would predict. We further suggest hybridization management actions should have clearly defined goals and be explicit about their assumptions. Finally, we provide an example of our framework applied to a common management action used to manage hybrid invasions: the use of barriers.

IDENTIFYING LAKE TROUT SPAWNING LOCATIONS IN SWAN LAKE, MONTANA

Benjamin S. Cox and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Montana State University, Bozeman, Montana 59717

Leo R. Rosenthal, Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, Montana 59901

Knowledge of spawning locations is critical to the management of invasive lake trout (*Salvelinus namaycush*) populations by providing areas to efficiently target sexually mature fish. In Swan Lake, Montana spawning locations were identified using acoustic telemetry, short-set gill nets, and in-situ egg nets. Telemetry locations were recorded manually after dark with a directional hydrophone from mid-October to mid-November in 2007 and 2008. Two primary sites were identified as potential spawning areas based on kernel-density analysis of 2007 telemetry locations. In 2008, 30 in-situ egg nets were buried by SCUBA divers at each of the 2007 locations to confirm egg deposition. One gill net (274.2 m long X 2.4 m deep with 5.08-cm bar mesh) was set at each primary spawning location once weekly through the tracking period in 2008 to confirm the presence of ripe lake trout and to explore the efficacy of targeting adult lake trout for removal at these locations. Kernel-density analysis of telemetry data identified the same primary spawning locations in 2008. Catch per unit effort of adult lake trout was > 4 times greater at these locations versus lake-wide gill netting. Egg density varied from 38 eggs/m² to 114 eggs/m² at the two spawning locations. Lake trout spawning locations were confirmed in Swan Lake, Montana. We recommend targeting these areas as an effective option for removing adult fish from the population.

LAKE TROUT REMOVAL EFFORTS CONTINUE IN YELLOWSTONE LAKE

Philip D. Doepke, Patricia E. Bigelow, Brian D. Ertel, and Todd M. Koel, Aquatic Sciences Section, Yellowstone National Park, P.O. Box 168, Mammoth Hot Springs, WY 82190

Lake trout (*Salvelinus namaycush*) removal efforts in Yellowstone lake began in 1995, the year following documentation of this non-native fish in Yellowstone Lake. The objective of the removal program was to reduce lake trout to the point where they have a negligible effect on native Yellowstone cutthroat trout (*Oncorhynchus clarkii bouvieri*). Lake trout capture techniques have focused on mechanical, mostly gill netting, and more recently electrical methods. The lake trout infestation in Yellowstone Lake continues while fishery managers remove greater numbers of lake trout every year, 76,140 lake trout were removed in 2008, and 348,794 have been removed since 1995. Amount of gill net effort has increased every year with the exception of 2008, total effort is mainly dependent on ice-off dates, and seasonal crew staffing and experience. Removal strategies target both adult and juvenile components of the population. Catch per effort of juvenile lake trout has steadily increased since 2002

with 2008 being the highest on record (5.0 lake trout/100 m gill net set/night) since 1998. However, cutthroat trout have not shown a positive response to lake trout removal efforts. Numbers of spawning cutthroat trout in Clear Creek, a tributary to Yellowstone Lake, are at the lowest levels ever recorded with just 538 counted. Yellowstone cutthroat trout gill net assessment, which averaged over 15 fish/net in 1994, has averaged ~ 9 fish/net over the past 5 yrs. Continued suppression of lake trout is imperative for there to be a healthy Yellowstone Cutthroat trout population in Yellowstone Lake.

RESTORING THE BALANCE: THE FISH & WILDLIFE PROGRAMS AT THE CRESTON FISH & WILDLIFE CENTER

Don Edsall, Rox Rogers, and Bob Lee, USDI Fish and Wildlife Service, Creston Fish and Wildlife Center, 780 Creston Hatchery Road, Kalispell, MT 59901

The Creston Fish and Wildlife Center is a USDI Fish and Wildlife Service facility located in Creston, Montana. There are three FWS programs working from the Center; the Creston National Fish Hatchery, Ecological Services and Partners for Fish and Wildlife. These programs work with partners and stakeholders to protect and conserve wildlife habitat, and to restore and enhance the fish and wildlife species who occupy these healthy habitats. This poster provides an introduction to the programs at the Center and a sampling of the species they strive to preserve.

ARCTIC GRAYLING EMERGENCE AND DEVELOPMENT IN ODELL CREEK, RED ROCK LAKES NATIONAL WILDLIFE REFUGE, MONTANA

Rebekah Levine, Department of Earth and Planetary Sciences, University of New Mexico, MSC03-2040, 1 University of New Mexico, Albuquerque, New Mexico 87131

Jeffrey M. Warren, Red Rock Lakes National Wildlife Refuge, 27820 Southside Centennial Road, Lima, Montana 59739

Glenn Boltz, USDI Fish and Wildlife Service, 4052 Bridger Canyon Road, Bozeman, Montana 59715

Red Rock Lakes National Wildlife Refuge in southwest Montana supports the last endemic population of adfluvial Arctic grayling (*Thymallus arcticus*) in the contiguous United States. The population has been declining for the last several decades and there is concern about its persistence on the Refuge. The goal of this study was to understand how Odell Creek is contributing to the maintenance of this critical population by investigating its use as a spawning stream and rearing ground for grayling fry as well as inventorying overall habitat. Surveys were conducted in 2006 and 2007 to determine the timing of spawning and emergence, to track fry movement in the creek and to make determinations about fry habitat selection. In 2006, 1868 fry were observed throughout the survey period with fry showing apparent movement downstream. During the same period in 2007 only 311 fry were observed. The majority were concentrated at the creek mouth. An a priori suite of models determined that there was little correlation between fry abundance and habitat. The most important parameter for fry abundance in 2006 was water velocity. Fry >30 mm were rarely observed; at this life stage they have improved swimming ability and are able to select different habitat types. Odell Creek's habitat was surveyed and determined to be a suitable grayling stream. The difference in fry abundance between years appears attributable to timing and intensity of

spring runoff which affects spawning and incubation periods. This indicated the importance of climate as a factor affecting this population.

ARE RESERVOIRS ECOLOGICAL SINKS FOR RECRUITMENT OF PALLID STURGEON?

Steven H. Ranney and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Montana State University, P.O. Box 173460, Bozeman, Montana 59717

Molly A. H. Webb and Kevin M. Kappenman, USDI. Fish and Wildlife Service, Bozeman Fish Technology Center, 4050 Bridger Canyon Road, Bozeman, Montana 59715

William M. Gardner, Montana Fish, Wildlife and Parks, 215 W. Aztec Drive, PO Box 938, Lewistown, Montana 59457

Natural recruitment of pallid sturgeon (*Scaphirhynchus albus*) in the Missouri River upstream of Fort Peck Reservoir has been nonexistent for at least four decades. Deviations from the natural river morphology are likely driving the lack of natural recruitment. An obligate riverine species, pallid sturgeon require large drift distances (> 245 km) before transitioning to benthic habitats. Insufficient lotic habitat downstream from hatch locations because of river impoundment is one hypothesis for the lack of pallid sturgeon recruitment. During 2008, we sampled the aquatic habitat in the headwaters of Fort Peck Reservoir. The headwaters were described as separate habitat units based on visible characteristics, i.e., “river,” “transition zone,” “side channel,” “lake,” and “reference.” Velocity decreased longitudinally through the headwaters of Fort Peck Reservoir suggesting that there is insufficient drift distance for pallid sturgeon to transition through the larval stage. Water-quality data examined at the diel scale indicated abrupt changes in temperature, dissolved oxygen, and pH that may negatively influence larval pallid sturgeon. These diel fluctuations in water quality parameters correlate with an increase in discharge from Canyon Ferry Reservoir. Further research includes spatial analysis of larval drift dynamics, evaluating larval sturgeon survival under fluctuating water-quality conditions, and experimentally evaluating larval pallid sturgeon survival in the headwaters environment.

MOVEMENT AND SPAWNING LOCATIONS OF SHOVELNOSE STURGEON AND PALLID STURGEON IN THE MISSOURI RIVER ABOVE FORT PECK RESERVOIR

Ryan R. Richards and Christopher S. Guy, U.S. Geological Survey, Montana Cooperative Fishery Research Unit, Montana State University, Bozeman, MT 59717

Susan L. Camp, USDI Bureau of Reclamation, Montana Area Office, P.O. Box 30137, Billings, Montana 59107

William M. Gardner, Montana Department of Fish, Wildlife and Parks, P.O. Box 938, 2358 Airport Road, Lewistown, MT 59457

The lack of recruitment by pallid sturgeon (*Scaphirhynchus albus*) over that last 40 yrs in the Missouri River above Fort Peck Reservoir has caused their abundance to decrease to less than 150 individuals. Interestingly, shovelnose sturgeon (*Scaphirhynchus platyrhynchus*) exhibit recruitment in the same area and their abundance is higher than most other rivers in the U.S. Understanding the dichotomy between the two species with respect to recruitment is

receiving much attention throughout the Mississippi River basin. The objectives of this study are to identify spawning locations and the effects of varying discharge on spawning locations and spawning movements for pallid sturgeon and shovelnose sturgeon. Two pallid sturgeon and 39 shovelnose sturgeon were radio tagged and tracked from 1 May 2008 to 5 July 2008. Unfortunately, no data is available for pallid sturgeon movement due to small sample size and loss of one of the radio-tagged fish mid-study. Seventy-four percent of the shovelnose sturgeon moved downstream during the spawning period. Of those that moved downstream the average movement was 39 km. One fish moved 123 km downstream and this was the largest movement of any fish tracked. Only 26 percent of the shovelnose sturgeon moved upstream and their average movement was 51.5 km. Shovelnose sturgeon concentrated in two areas during the spawning period; river kilometer 3090 to 3190 and 3235 to 3270. These data will provide a greater understanding of sturgeon (*Scaphirhynchus* spp). spawning movements and spawning locations in the upper Missouri River and will help guide management decisions aimed at recovery of pallid sturgeon.

LARGE WOODY DEBRIS DEPLETION RATES IN WESTERN MONTANA STREAMS

Brian Sugden, Plum Creek Timber Company, Inc., P.O. Box 1990, Columbia Falls, Montana 59912

Ron Steiner, Plum Creek Timber Company, Inc., P.O. Box 1990 , Columbia Falls, Montana 59912

Maintenance of in-stream habitat diversity through recruitment of large woody debris (LWD) is a principal goal of Plum Creek Timber Company's Native Fish Habitat Conservation Plan (NFHCP). Because LWD loads are governed by long-term processes such as riparian tree growth and mortality, direct measurement of LWD to judge effectiveness of management actions is unlikely to provide timely feedback for adaptive management. Instead, trends must be forecast using models. For the NFHCP, the Riparian Aquatic Interaction Simulator (RAIS, Welty et al. 2002) was used to support development of riparian management options. To help validate this model, several key assumptions are being field-tested as part of the NFHCP's adaptive management program. This study examines the effects of channel size and gradient on annual in-stream LWD depletion rates. Thirty-one sample sites were randomly selected from among a pool of stream segments that represent combinations of channel types and sizes for perennial streams on Plum Creek lands. At each 100-m site, all pieces of LWD with minimum qualifying dimensions of 10-cm diameter and 2-m length in zones 1-3 (Robison and Beschta 1990) were tallied and marked with numbered aluminum tags. Resurveys of these sites were completed in 2007. This poster summarizes five-year LWD depletion rates, and the factors most associated with LWD movement.

MONTANA UNAUTHORIZED FISH INTRODUCTIONS

Jim Vashro, Regional Fisheries Manager, Montana Fish, Wildlife & Parks, 490 N. Meridian, Kalispell, MT 59901

Poster display will consist of a map of Montana with illegal fish introduction locations marked by symbols. Display would include a list of species and distribution by region. A copy of the database would be available with an invitation for fisheries personnel to review their area and report any fish introductions not yet documented. Report forms would be available.

BIOLOGICAL SCIENCES – TERRESTRIAL

VIABILITY OF USER-PAID SYSTEM OF WILDLIFE CONSERVATION IN MONTANA

Thomas R. Baumeister, Jeffrey T. Herbert, Mike Lewis, Justin A. Gude, Montana Fish, Wildlife, and Parks, Helena, Montana 59620

Julie A. Cunningham, Montana Fish, Wildlife, and Parks, Bozeman, Montana 59718

Montana Fish, Wildlife and Parks (FWP) is entrusted with the responsibility to conserve all species of fish and wildlife to meet a variety of public interests and values. To date, users of these resources, namely hunters and anglers, have provided the financial and political support for the state to achieve this mandate. With this presentation, we make the business case for why user participation is critical to the future of fish and wildlife stewardship in the state, and compare current participation trends in Montana to those of other states in anticipation of what we might experience in the future. This will be followed by a brief discussion of the primary factors leading to changes in participation, and how they relate to the user-paid model of wildlife conservation using a conceptual framework. We will then present and discuss a variety of strategic initiatives Montana could explore to address the implications of a changing user demographic. We will conclude by providing an overview of ongoing efforts by FWP to date including the role of marketing and branding of our services and products. We're hopeful that these two presentations will be supportive of TWS-Montana Chapter's efforts to explore ways in which we can promote sound stewardship of wildlife and their habitats in Montana in a manner that is economically and socially sustainable.

WOLF PREY PREFERENCES IN MULTIPLE PREY SYSTEMS: INSIGHTS FROM THE MADISON HEADWATERS OF YELLOWSTONE NATIONAL PARK

Matthew S. Becker, and Robert A. Garrott, and Claire N. Gower, Ecology Department, Montana State University, Bozeman, Montana 59717

Patrick J. White, Yellowstone Center for Resources, Yellowstone National Park, Wyoming 82190

We studied wolf prey selection and kill rates during 1996-1997 through 2006-20007 winters in a newly established one predator two-prey system in central Yellowstone National Park. Prey differed substantially in their vulnerability to wolf (*Canis lupus*) predation and wolves preyed primarily on elk (*Cervus elaphus*) but also used bison (*Bison bison*) to varying degrees within and among winters. Winter severity, wolf abundance, distribution, and prey selection varied during the study, concurrent with variations in the demography, distribution, and behavior of elk and bison. Patterns of prey selection trends were strongly correlated to elk calf abundance. While wolves increasingly killed bison with increasing bison: elk ratios, snow pack duration, and wolf numbers, they did not appear to change their preference for elk. Similarly, variation in elk kill rates were not related to or reduced by increases in bison kill rates. The wolf functional response for elk was a Type II, indicative of a preferred prey, and strongly influenced by wolf abundance, as it was positively correlated with increased competition and anti-predator responses of elk. Prey-switching evaluations indicated increasing selection of bison with increasing bison: elk ratios, however no concurrent

decrease in elk predation occurred. Increased bison predation is not solely dependent on relative abundance of the two prey species; therefore it is unlikely at this time that wolf prey-switching will stabilize the system. The pervasive influence of differential vulnerability among prey species and age classes and its effects on the potential trajectories of wolf-ungulate systems in Montana is discussed.

GIS-BASED TOOLS TO IMPROVE LAND USE PLANNING FOR WILDLIFE CONSERVATION

Brent L. Brock and Lance Craighead, The Craighead Center for Landscape Conservation, Craighead Environmental Research Institute, 201 South Wallace Ave., Suite B2D, Bozeman, Montana 59715

Conserving wildlife and habitat connectivity in the face of a growing human population is one of the greatest challenges facing wildlife managers in the 21st Century. The Northern Rocky Mountain region is experiencing some of the most rapid human population increases in the United States, and rural sprawl is now recognized as one of the most serious threats facing Montana wildlife in the near future. To address this challenge, land use planners must incorporate the best available science about wildlife requirements into planning decisions and policy. We developed a suite of GIS-based tools to simplify incorporating scientific information into landscape-level land use planning for wildlife. These tools are designed to be flexible to accommodate a range of wildlife conservation objectives. The current suite includes tools for estimating appropriate development densities likely to provide adequate habitat or movement connectivity for specified wildlife targets, as well as evaluate existing landscapes, or potential development scenarios, to estimate their potential for supporting wildlife. These tools were employed in conjunction with the development of a wildlife overlay in the Madison Valley, Montana. The Madison Valley provides an example of how these tools can assist with creating criteria for development near or within important wildlife areas.

WINTER DISTRIBUTION, HABITAT USE, AND BROWSE UTILIZATION PATTERNS OF THE SHIRAS MOOSE ON THE MOUNT HAGGIN WILDLIFE MANAGEMENT AREA

Braden O. Burkholder and Vanna J. Boccadori, Montana Fish, Wildlife and Parks, Butte, Montana 59701

Robert A. Garrott, Fish and Wildlife Management Program, Department of Ecology, Montana State University, Bozeman, Montana 59717

Moose (*Alces alces shirasi*) populations across Montana have expanded in the last century, both in geographic range and in population size. This expansion has had a negative impact on moose winter range in some locations where moose have overutilized key browse species such as aspen and willow. Excessive and unsustainable browsing has the potential to reduce local biodiversity and carrying capacity of moose and other ungulates. The browse species of interest in this study are willow (*Salix* spp.), a highly palatable and abundant browse source for moose on many winter ranges, including our study area in southwestern Montana. Knowledge of spatial and temporal patterns of moose willow community use and willow utilization patterns is limited in Montana and would be helpful in moose population

management. The objectives of this study were to determine patterns of willow community use by selected female moose during winter and to quantify willow utilization across the study area to examine population scale habitat use through browse patterns. To accomplish these objectives we deployed GPS collars on 12 cow moose in the winters of 2007 and 2008 and completed large scale, systematic browse surveys in the spring of 2008. Preliminary results indicated that cow moose spend the majority of the winter in or adjacent to willow communities, but overall willow utilization across the study area was low. Our data suggested that while moose have the potential to significantly impact willow communities, this did not appear to be the case on the Mount Haggin WMA at current moose densities.

PUBLIC OPINION AND KNOWLEDGE OF GRIZZLY BEARS IN THE CABINET-YAAK ECOSYSTEM

Sarah Canepa, Troy, Montana 59935

Kimberly M. Annis*, Montana Fish, Wildlife and Parks, Libby, Montana 59923

Wayne Kasworm, USDI Fish and Wildlife Service, Libby, Montana 59923

To measure the public's understanding of grizzly bears and their management in the Cabinet-Yaak Ecosystem (CYE), a telephone survey was conducted in Lincoln and Sanders County, Montana. In the summer of 2007, 502 residents of the seven communities within the CYE answered questions about their knowledge and opinions of grizzly bears (*Ursus arctos*) in the CYE and of related management activities. Ninety percent of respondents felt that humans can prevent most conflicts with grizzly bears and 62 percent stated that they would accept changes to current garbage disposal methods if it would help prevent problems with grizzly bears. Fifty-seven percent supported the recovery of the grizzly bear population in the CYE. Support decreased to 44 percent for achieving a population goal of 100 bears. Support increased to 75 percent if population recovery could be achieved without using augmentation. Thirty-three percent stated they were unaware of road access restrictions on National Forest lands, due in part to grizzly bear recovery efforts. The majority of respondents indicated some level of support for grizzly bears, yet had concerns over specific management actions used to achieve population recovery. Respondents were more aware of augmentation efforts in the early 1990's than of more recent efforts, suggesting that managers need to keep the public better informed. Educational efforts may benefit residents' understanding of general grizzly bear biology and of related management practices.

MONTANA'S CONSERVATION STRATEGY FOR ROCKY MOUNTAIN BIGHORN SHEEP

Tom Carlsen, Montana Fish, Wildlife and Parks, Townsend, Montana 59644

Montana Fish, Wildlife and Parks (FWP) is currently in the process of developing a comprehensive Conservation Strategy for Rocky Mountain bighorn sheep (*Ovis canadensis*). The strategy includes the history of bighorn sheep in Montana from decline to recovery. Direction on how FWP monitors and manages populations, herd health, and bighorn habitat is defined. Protocols for resolving situations where bighorn sheep and domestic sheep/goats commingle, recommendations regarding use of domestic sheep/goats for noxious weed control, and a protocol for responding to die-offs have been developed. A Translocation program, including processes for identifying and evaluating potential habitats and prioritizing

transplant sites are included in the strategy. Integral aspects of the strategy are narratives for each hunting district or population. These narratives include a complete history of the individual population, overall management goals, and objectives for habitat, access, and population demographics. As part of the strategy, how individual populations are managed through hunting and translocation is clearly linked to monitoring efforts.

GRIZZLY BEAR TREND MONITORING RESEARCH IN THE NCDE: AN UPDATE, 2004-2008

Tonya Chilton and Richard Mace, Montana Fish, Wildlife, & Parks, Kalispell, Montana 59901

In 2004, monitoring of grizzly bear (*Ursus arctos*) population trend in the Northern Continental Divide Ecosystem (NCDE) was initiated by Montana Fish, Wildlife and Parks (MFWP) in cooperation with other state and federal agencies. Also in 2004 an interagency field effort was undertaken to enumerate population size of grizzly bears over the entire NCDE using DNA methods. From these data, researchers have recently estimated population size with confidence limits and relative bear density across the NCDE. However, Managers and the public agree that a companion program is needed that would track population trend and female vital rates over time and provide ancillary information on other indices of population health. Having estimates of both size and trend will greatly improve our collective knowledge of grizzly bear ecology and provide more measurable and precise information with which to judge the status of the grizzly population in the NCDE. The purpose of this long-term trend program is to monitor the vital population parameters of grizzly bears by assessing the survival and reproductive rates, as well as trend, by following a number of radio-collared female grizzly bears. We summarize the objectives and field design of population trend and report on our success to date.

VIABILITY OF RESIDENT DEER AND ELK HUNTER PARTICIPATION IN MONTANA

Julie A. Cunningham, Montana Fish, Wildlife, and Parks, Bozeman, Montana 59718

Justin A. Gude, Thomas R. Baumeister, and Jeffrey T. Herbert, Montana Fish, Wildlife, and Parks, Helena, Montana 59620

The majority of funding for Montana FWP programs is generated or tied to sales of deer (*Odocoileus* spp.) and elk (*Cervus elaphus*) hunting licenses. Future declines in deer or elk hunter participation could therefore have negative consequences for wildlife management and conservation programs. To investigate deer and elk hunter participation rates in Montana, we made use of the FWP Automated Licensing System (ALS) database. Since 2002, the number of individuals buying deer and elk licenses has been roughly stable. We converted the data in ALS to a 6 (yrs) by 255,851 (unique individuals) matrix of resident hunter encounter histories for the period 2002-2007. Using a multi-state mark recapture model, we estimated hunter retention rates, hunter license buying probabilities, and transition probabilities between major age classes, considering hunter sex and region of residence as group covariates. We then used FWP hunter education databases to estimate hunter recruitment rates. Using these estimated rates, we parameterized and analyzed a stage-based population projection matrix. Estimated recruitment rates for teenagers following hunter education courses were sufficient to stabilize

the trend in deer and elk hunter participation, given our estimates of retention and license buying rates, in agreement with the overall trend in license sales for 2002-2007. Based on sensitivity and elasticity analyses of the projection matrix, future hunter participation rates and license sales would be most influenced by increases in recruitment rates and license buying probabilities for middle-aged adult and teenage males. We discuss the implications of these results for FWP hunter recruitment and retention efforts and social trends.

NUTRIENT ALLOCATION IN EGG FORMATION OF FEMALE LESSER SCAUP ON LOWER RED ROCK LAKE, RED ROCK LAKES NATIONAL WILDLIFE REFUGE

Kyle A. Cutting and Jay J. Rotella, Montana State University, Department of Ecology, Bozeman, Montana 59717

Jeffrey M. Warren, USDI Fish and Wildlife Service, Red Rock Lakes National Wildlife Refuge, Lima, Montana 59739

Susan E. Wainwright and John Y. Takekawa, U.S. Geological Survey, San Francisco Bay Estuary Field Station, Vallejo, California 94592

North American lesser scaup (*Aythya affinis*) populations have been declining for nearly three decades. Recent evidence suggests that decreases in the quantity and quality of wintering and spring staging habitat may be reducing the ability of scaup to obtain the dietary resources necessary to undertake breeding activities. We used stable isotope techniques to explore the relative contributions of endogenous (body) and exogenous (local) reserves in clutch formation and how nutrient contributions changed throughout the clutch formation period. Female scaup body tissue samples (blood, claw, and lipid) were collected from mid-May (shortly after their arrival on the study site) through the nesting period during 2007-08; eggs were collected from nests during 2006-2008. Tissue isotopic signatures from mid-May were significantly enriched in stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotope values compared to local dietary sources. Eventually tissue signatures equilibrated with local dietary resources by the end of the egg laying period in early July. Egg albumen, and to a lesser extent lipid-free yolk protein, were primarily derived from local sources. However, approximately half of yolk lipid was derived from endogenous body reserves. There was no significant relationship between isotopic signatures of eggs and the laying date. Our results indicate that while local resources are very important for meeting the nutrient demands of clutch formation, a majority of lipids and some protein are obtained from the spring staging areas.

BLOOD-LEAD LEVELS OF FALL MIGRANT GOLDEN EAGLES IN WEST-CENTRAL MONTANA

Robert Domenech, Raptor View Research Institute, Missoula, Montana 59801

Heiko Langner, Environmental Biogeochemistry Laboratory, University of Montana, Missoula, Montana 59812

Lead has long been documented as a serious environmental hazard to eagles and other predatory opportunistic and scavenging avian species. Due to lead poisoning in the Bald Eagle (*Haliaeetus leucocephalus*) the use of lead shot for waterfowl hunting on federal and state lands was banned in 1991. More recently, lead poisoning from spent ammunition

has been identified as the leading cause of death in California Condors (*Gymnogyps californianus*), prompting the recent ban of lead ammunition within the “California Condor Recovery Zone.” Another study on Common Ravens (*Corvus corax*) in Wyoming has shown a direct correlation between elevated blood-lead levels and the on-set of rifle hunting season. Indeed, there is overwhelming evidence showing that lead is still prevalent in the environment and mounting data points to lead based rifle bullets as the primary source. We sampled blood from 42 Golden Eagles (*Aquila chrysaetos*) captured on migration during the fall of 2006 and 2007 to quantify a suite of possible heavy metal contaminants, with an emphasis on lead. Lead was measured in micrograms per deciliter (ug/dl) and ranged from 0 - 481 ug/dl. The blood-lead levels were broken in four exposure stages and our results were as follows: eagles with 0 -10 ug/dl ($N = 18$) were considered background, 10 – 60 ug/dl ($N = 19$) sub-clinically exposed, 60– 00 ug/dl ($N = 2$) clinically exposed and any eagle with ≥ 100 ug/dl ($N = 3$) were considered acutely exposed. In all, we found that 58 percent of the 42 Golden Eagles sampled had elevated blood-lead levels.

A COOPERATIVE APPROACH TO ELK MANAGEMENT IN THE WILDLAND/URBAN INTERFACE OF MISSOULA, MONTANA - A DYNAMIC STRATEGY FOR A GROWING PROBLEM

Victoria L. Edwards, Montana Fish, Wildlife and Parks, Missoula, Montana 59804

Shawn M. Cleveland,* Wildlife Biology Program, University of Montana, Missoula, Montana 59812

Bert Lindler,* North Hills Homeowner, Missoula, Montana 59808

The Missoula Valley in western Montana is home to nearly 800 wintering elk (*Cervus elaphus*), of which 300 are from the North Hills elk herd. From 1980 to 2007, this herd grew from 17 to 290 elk with a 48-percent growth rate occurring between 2000 and 2007. Without an effective harvest, this population is projected to double every five years. With increased residential development in elk winter range, a diverse public’s opinion on the management objectives of the herd, and the herd’s juxtaposition to the City of Missoula and the Rattlesnake Wilderness and National Recreation Area, wildlife biologists have needed to become more creative with their management strategies. To more effectively manage the herd at a sustainable level and to keep it wild, wildlife biologists from Montana Fish, Wildlife and Parks have coordinated and collaborated with numerous landowners and staff at the University of Montana and the USDA Forest Service. This presentation is an integration of the social and biological sciences, with discussions on the successes and failures of tested strategies to manage elk in the wildland/urban interface. The discussion will include the perspective and efforts of a North Hills homeowner, and data from a Master student’s thesis project on survival, habitat use, daily/seasonal movements, and elk redistribution related to hunting pressure as collected with 10 GPS and 11 VHF collars. The discussion will conclude with a description of the adaptive management approaches utilized by Montana Fish, Wildlife and Parks’ wildlife biologists, and the efficacy of those strategies.

EFFECTS OF WOLF PREDATION ON THE MADISON HEADWATERS ELK HERD: INSIGHTS FOR ELK AND WOLF MANAGEMENT IN MONTANA

Robert A. Garrott, Ecology Department, Montana State University, Bozeman, Montana 59717

Patrick J. White, Yellowstone Center for Resources, Yellowstone National Park, Wyoming 82190

Claire N. Gower and Matthew S. Becker, Ecology Department, Montana State University, Bozeman, Montana 59717

Kenneth L. Hamlin, Montana Fish Wildlife and Parks, Bozeman, Montana 59717

Studies of the non-migratory Madison Headwaters elk (*Cervus elaphus*) herd in Yellowstone National Park revealed that the herd appeared to be regulated near ecological carrying capacity by food limitation prior to the reestablishment of wolves (*Canis lupus*). Eleven years of post-wolf data indicated a substantial proportion of wolf predation was additive and overwhelmed any potential for the elk to demographically compensate. Thus, wolf predation resulted in a dramatic decrease in elk abundance and the system transitioned from being bottom-up regulated in the absence of a significant predator to strong top-down limitation due to wolf predation. It is uncertain if predation will ultimately regulate the elk population at a lower, alternate state or if predation and other factors influencing elk vulnerability will interact to result in further decreases in elk abundance. Fundamental to this question is the role of alternative prey and the interactions of winter severity and landscape heterogeneity on the vulnerability of elk to wolf predation. We contrast the impacts of wolf predation on elk in different drainages of the Yellowstone study area, as well as nearby areas studied outside the Park, to gain insights into the varying impacts of wolves on elk populations that occupy diverse landscapes in Montana. We suggest that wolves will have little to modest impacts on numbers of elk in most areas of the state, but that additive wolf predation could result in reduced elk numbers in some herds that winter in deep snow, forested environments where limited conflicts with livestock production can result in higher wolf: elk ratios.

ELK BEHAVIORAL RESPONSES TO THE REESTABLISHMENT OF WOLVES: INTEGRATING MULTIPLE STRATEGIES TO ACCOMMODATE COMPETING DEMANDS

Claire N. Gower, Robert A. Garrott, and Matthew S. Becker, Fish and Wildlife Management Program, Department of Ecology, Montana State University, Bozeman, Montana 59717

P.J. White, Yellowstone Center for Resources, Yellowstone National Park, Wyoming, 82190

Over the past few decades a large body of literature has provided evidence that predators can influence the ways in which prey behave. This in turn may influence prey demography and predator-prey dynamics, and therefore may influence prey populations independent of direct killing. Using data collected from 1991 to 2007, we evaluated the behavior of elk (*Cervus elaphus*) in the Madison headwaters area of Yellowstone National Park in response to the colonization and establishment of wolves (*Canis lupus*). Changes in home range size, fidelity, group size, foraging, and large-scale spatial responses were evaluated. Prior to wolf colonization, grouping behavior was relatively stable and predictable as elk attempted to conserve energy and decrease starvation risk in the absence of wolves. Following reintroduction group size and group size variation increased. This more dynamic behavior

likely reflects a strategy to minimize predation and maximize food acquisition. After wolf colonization, elk moved more over the landscape as they were increasingly encountered, attacked, and displaced by wolves. Home ranges were slightly larger, with some decreases in fidelity. Long-distance dispersal and migratory movements were also adopted away from high-density wolf areas. These apparent predator-avoidance movements were never observed prior to wolf colonization or from areas where predation risk was lower. The decision to forage was heavily influenced by local snow, habitat, and time of day but remained relatively stable with and without the presence of wolves. We suspect that this lack of any substantial change in foraging behavior illustrates that elk can maintain the same level of foraging time and retain a relatively constant level of nutrition. Together these results suggest that in a harsh winter environment, elk can adaptively manage their behavior to cope with environmental constraints both in the presence and absence of wolves.

GRIZZLY BEAR POPULATION AUGMENTATION IN THE CABINET MOUNTAINS OF NORTHWEST MONTANA

Wayne Kasworm, USDI Fish and Wildlife Service, Libby, Montana 59923

Kimberly M. Annis, Montana Fish, Wildlife and Parks, Libby, Montana 59923

Timothy Manley, Heather Reich, Derek Reich, and Jim Williams, Montana Fish, Wildlife and Parks, Kalispell, Montana 59901

Chris Servheen, USDI Fish and Wildlife Service, Missoula, Montana 59812

The Cabinet Mountains grizzly bear (*Ursus arctos*) population was estimated at 15 or fewer individuals in 1988 and believed to be declining toward extinction. In response to this decline, a test of population augmentation techniques was conducted during 1990-1994 when 4 subadult female grizzly bears were transplanted into the area. Two criteria were identified as measures of success: bears must remain in the target area for one year, and bears should ultimately breed with native male grizzly bears and reproduce. Three of four bears remained in the target area for one year or more. One bear died after one year. Reproductive success of any of the remaining individuals could not be established until 2006 when genetic analysis of hair snag samples collected from 2002-2005 was completed. This analysis indicated that at least one of the transplanted bears remained in the Cabinet Mountains and had reproduced. The detected bear was transplanted in 1993 as a 2-year-old and was identified by a hair snag within 5 mi of the original release site. Genetic analysis indicated she had produced at least 4 offspring and those offspring had also reproduced. This information indicates that the original test of augmentation was successful with at least one of the transplanted individuals. The success of the grizzly bear augmentation test has prompted continuation of this effort. The Northern Continental Divide Ecosystem area of north central Montana has been the source of four additional bears transplanted to the Cabinet Mountains during 2005-2008. Monitoring of these recent transplants is described.

DEMOGRAPHY AND GENETIC STRUCTURE OF A RECOVERING GRIZZLY BEAR POPULATION

Katherine C. Kendall, U. S. Geological Survey–Northern Rocky Mountain Science Center, Glacier Field Station, Glacier National Park, West Glacier, Montana 59936

Jeffrey B. Stetz and Amy C. Macleod, University of Montana Cooperative Ecosystem Studies Unit, Glacier Field Station, Glacier National Park, West Glacier, Montana 59936

John B. Boulanger, Integrated Ecological Research, Nelson, BC V1L 4L4, Canada

David Paetkau, Wildlife Genetics International, Nelson, BC V1L 5P9, Canada

Gary C. White, Department of Fish, Wildlife and Conservation Biology, Colorado State University, Fort Collins, Colorado 80523

The threatened grizzly bear (*Ursus arctos*) population in northwestern Montana has been managed for recovery since 1975, yet no rigorous data were available to monitor program success. We assessed population status using data from one of the world's largest noninvasive genetic sampling efforts and 33-years of physical captures. Our population estimate, $N = 765$ ($CV = 3.8\%$) was double the working estimate. Based on our results, the recent human-caused mortality rate approached a sustainable 4 percent although the high proportion of female mortalities raises concern. Genetic interchange has recently increased in areas exhibiting generations of low gene flow. This study illustrates the power of molecular techniques to rapidly assess populations at landscape scales and provide detailed demographic and genetic data needed to guide and evaluate recovery efforts.

RELATIONSHIPS BETWEEN ELK AND NONNATIVE WEEDS ON MONTANE WINTER RANGES IN WESTERN MONTANA

Michel T. Kohl, Mark Hebblewhite, and Shawn M. Cleveland, Wildlife Biology Department, University of Montana, Missoula, Montana 59801

Through physiological and competitive mechanisms, spotted knapweed has become established in many critical winter-range habitats, primarily in the northwestern United States and southwestern Canada. In areas of high density, elk (*Cervus elaphus*) may reduce the competitive ability of native grasslands through overgrazing. I predict that the benefits of weed spraying to reduce knapweed biomass are reduced in areas of high elk density and elk density and knapweed cover are positively correlated with high elk density creating high levels of weed biomass. A linear regression was calculated using vegetation data against kernel density estimates obtained from GPS telemetry locations in the North Hills Elk Herd of Missoula Valley, Montana. This analysis shows a trend for increased and decreased percent cover of knapweed and native grasses at high elk densities respectively, providing support for our overgrazing hypothesis through the mechanism of apparent competition. Fecal diet analysis further shows avoidance of knapweed and a selection for native grasses and forbs as well as invasive cheat grass. Finally, elk exclosures were constructed and will be measured annually for comparisons between forage class (native, invasive) biomass and elk densities on herbicide treatment areas using a statistical analysis of covariance. This information is essential in understanding the secondary effects of weed spraying in areas of high ungulate densities. With this understanding, management may be better informed how to spend limited resources for invasive weed control. If benefits of weed spraying are diminished at high ungulate densities, managers will need to adopt very different weed control treatments.

FROM MAGIC TO TRAGIC: THE HISTORY OF WOLF RECOVERY AND MANAGEMENT IN NORTHWEST MONTANA

Kent Laudon, Montana Fish Wildlife and Parks, Kalispell, Montana 59901

In 1973, Northwest Montana wolf (*Canis lupus*) recovery began with Dr. Bob Ream, University of Montana Wolf Ecology Project, and a handful of volunteers trying to verify presence. Verified presence increased slowly until 1986 when the first pack and reproduction, the Magic Pack, was documented. Since then and until recently, the Northwest Montana Recovery Area (NMRA) was the slowest growing, and survivorship of individual wolves was the lowest of the three recovery areas within the Northern Rocky Mountain Population (NRMP). In the last three years the NMRA population rate of increase is now similar to the Greater Yellowstone Recovery area. The NRMP wolf population reached recovery goals in 2002 and will soon be delisted. During this 35 year recovery period numerous biologists working for University, Federal, State, Tribal, and private interests monitored and managed the population and devoted much to the success of the project. The numerous events, stories, and people behind its ultimate success are colorful and scattered with both accomplishments and set backs. Managers developed a complex bag of tools for population monitoring and management, livestock conflict resolution, and public relations unique to wolves. Wolf management is one of the more controversial programs that face wildlife professionals, and as the wolf population increases so do the challenges. An increased wolf population is met with increased public concerns on multiple issues including ungulate hunting opportunity, livestock interests, human safety, and domestic dog conflicts. Emotions have run high recently when managers have been severely criticized by diametrically opposing publics for either killing too many wolves or for white-tailed deer declines believed to be caused by wolves. The future holds yet more challenges with litigation, pending wolf hunts, as well as the same challenges of the past.

WHY IS THIS GRIZZLY BEAR IN MY BACKYARD? MANAGING HUMAN/GRIZZLY BEAR CONFLICTS IN NORTHWEST MONTANA

Timothy L. Manley, Montana Fish, Wildlife and Parks, Kalispell, Montana 59912

The question may seem simple, “Why is this grizzly bear (*Ursus arctos*) in my backyard”? A reasonable answer would be “Because it has probably found something to eat.” While the question and answer are basic, the solutions and prevention of future conflicts are more complex. From 1993 through 2008, there were 244 management captures of grizzly bears in Northwest Montana. These captures ranged from responses to grizzly bears causing property damage to incidental captures. Interagency Grizzly Bear Committee (IGBC) Guidelines give direction on capture and fate of the bear, but it is up to the Bear Conflict Management Specialists to work with both the grizzly bear and the people on the ground. Techniques to deal with the grizzly bears include relocation, on-site releases, hazing, aversive-conditioning, securing attractants, and removal from the population. Techniques to deal with the people include education, assistance in securing attractants, and citations. Just like people, grizzly bears are all individuals, and what might work in one situation might not work in all situations. Methods for dealing with and preventing conflicts is evolving as new techniques and technology gives grizzly bear managers new tools and options. The ultimate goals are to keep people safe and to keep grizzly bears alive and in the wild.

AN ECOLOGICAL RISK ASSESSMENT OF WIND ENERGY DEVELOPMENT IN MONTANA

Brian H. Martin, Amy J. Pearson, Brad D. Bauer, The Nature Conservancy, Helena, Montana 59601

In 2008 the United States led the world in wind power generation, providing 35 percent of the nation's new electrical generating capacity. Montana ranks fifth among states for wind energy potential. Wind facilities are not stand-alone features—they cover vastly more area than the footprint of the turbines, requiring extensive road systems and transmission corridors. The challenge for wind energy development in Montana is to produce relatively clean energy that does not contribute to global climate change, while minimizing impacts to biodiversity. We have completed an ecological risk assessment at coarse and fine scales for Montana. We utilized a diverse assemblage of wildlife species of concern, selecting for those that research suggests would be the most susceptible to wind energy development. We estimate that in total about 17 million ac of available good-to-superb wind energy potential exists within Montana, and of that, at least 7.7 million ac have a high risk to potentially impact species of concern. We strongly suggest that these areas be avoided as locations for wind energy development, rather than considering mitigation approaches, as the lands identified are often critical habitat for multiple species.

MODELING PREDICTED DISTRIBUTION AND LANDSCAPE-LEVEL HABITAT SUITABILITY FOR MONTANA WILDLIFE SPECIES

Bryce A. Maxell, Montana Natural Heritage Program, University of Montana, Helena, Montana, 59620

Scott Story and Joy Ritter, Information Management Bureau, Montana Department of Fish, Wildlife, and Parks, Helena, Montana 59620

Models predicting spatial distribution and habitat suitability are critical for natural resource managers making decisions that impact species for which there is limited information. We are using presence-only data in conjunction with pseudo-absences in program Maxent to model the distribution and landscape-level habitat suitability for Montana wildlife species. Our primary goals are to produce (1) continuous statewide outputs as a tool to identify variables that limit species' distributions and areas that need field surveys; (2) binary outputs that can be used to create lists of predicted species for various administrative boundaries and, (3) outputs showing marginal, suitable, and optimal habitat classes at a local landscape-level. To date, models have identified scale dependent responses to environmental variables, opportunities to extend the known ranges of species, areas that support potentially isolated populations in need of conservation efforts, areas that are critical for maintaining landscape connectivity, areas that may provide the best habitat for reintroduction of species that have declined, and areas where exotic and nonindigenous species are most likely to become established. In general, inductively based Maxent models provide realistic depictions of species distributions when survey data is available for a region. However, deductive models will still be important for representing some species distributions in areas lacking survey effort. Models will be used for various planning efforts including Montana Fish, Wildlife, and Parks' Crucial Areas and Corridors Assessment. Model outputs can be obtained from the Montana Natural Heritage Program or the Montana Fish, Wildlife, and Parks' Information Management Bureau.

STATUS OF LENTIC BREEDING AMPHIBIANS AND AQUATIC REPTILES IN MONTANA

Bryce A. Maxell and Dave Ratz, Montana Natural Heritage Program, University of Montana, Helena, Montana 59620

P. Stephen Corn, U.S. Geological Survey Northern Rocky Mountain Science Center, Aldo Leopold Wilderness Research Institute, Missoula, Montana 59807

D. Grant Hokit, Department of Natural Sciences, Carroll College, Helena, Montana 59625

We developed a statewide inventory and monitoring scheme for lentic breeding amphibians and aquatic reptiles in Montana. We stratified sampling by 11 ecoregions and surveyed 6741 potential lentic sites on public lands between 2000 and 2008 within 429 randomly selected 12-digit hydrologic-unit-code watersheds. Surveys and associated incidental observations have resulted in over 11,400 species occurrence records to date. Watershed and site breeding rates of 10 amphibian species and occupancy rates of four aquatic reptile species support previously noted declines of the Western Toad (*Bufo boreas*) and Northern Leopard Frog (*Rana pipiens*) in western Montana. We used classification trees to examine patterns in rates resulting from different groupings of major habitat features that are able to be affected with management actions. Seven of the 10 amphibian species and the Common Gartersnake (*T. sirtalis*) were detected at significantly fewer sites when fish were detected. Presence of emergent vegetation was positively associated with the proportion of sites where breeding or occupancy was detected for all but one of the amphibian and reptile species examined and appeared to partially mitigate the presence of fish. Resource managers could enhance habitats for wetland herpetofauna by (1) creating new lentic sites on the landscape either directly or through the reintroduction and protection of beaver, (2) creating emergent vegetation at portions of existing sites that currently lack it via rotational fencing to temporarily exclude grazing, and (3) eliminating some introduced fish populations. All observations and survey locations, including digital photographs of sites surveyed are available at: <http://nhp.nris.state.mt.us/Tracker>.

THE IMPACT OF CLIMATE VARIATION ON COLUMBIA SPOTTED FROG SURVIVAL IN A HIGH MOUNTAIN ECOSYSTEM

Rebecca McCaffery, Wildlife Biology Program, University of Montana, Missoula, Montana 59812

Bryce Maxell, Montana Natural Heritage Program, Helena, Montana 59620

Amid growing concern over the impacts of long-term climate change, a fundamental challenge for wildlife biologists is determining how animal populations will respond to a changing climate. In amphibians, little research has addressed how climate variation may affect vital rates and population sizes. We evaluated the relationship between annual age- and sex-specific survival rates and local and global climate variables using a nine-year mark-recapture dataset of Columbia spotted frogs (*Rana luteiventris*) from the Bitterroot Mountains. Local climate variables included peak snowpack, winter length, summer length, and growing degree days. Global climate variables included the Southern Oscillation Index (SOI) and the North Pacific Oscillation (NPO). We estimated annual survival for four age classes: juveniles, subadult and adult females, and adult males. We found that survival in this population was best predicted by snowpack and winter length. In this model, an increase in snowpack resulted in a decrease in juvenile ($b = -0.083 \pm 0.007$), adult male ($b = -0.036 \pm 0.016$), and

adult female ($b = -0.037 \pm 0.015$) survival, and had no significant effect on subadult survival ($b = -0.011 \pm 0.027$). An increase in winter length led to a slight increase in survival, but only for juveniles and adult males. These results suggest that a warming climate with less severe winters might be good for montane frog populations. Survival is only one vital rate, however, and future work will determine the influence of climate on other vital rates such as growth and fecundity, as well as examine the contribution of intrinsic drivers to population variation.

HUNTING ACCESS MANAGEMENT ON PRIVATE LANDS IN MONTANA

Caitlin McCoy and Tara Teel, Human Dimensions of Natural Resources Department Colorado State University, Fort Collins, Colorado

Mike Lewis, Montana Fish, Wildlife and Parks, Helena, Montana 59620

In the summer of 2008, a study of hunting access on private lands in Montana was conducted cooperatively by Colorado State University, Montana Fish, Wildlife and Parks, and the Western Association of Fish and Wildlife Agencies. The purpose of this effort was to examine the extent to which Montana landowners are providing hunting access and the approaches they use to manage access on their property. The focus was on access for hunting of deer, elk, antelope, and upland game birds (defined to include pheasant, sharp-tailed grouse, and Hungarian partridge). The study also included an assessment of landowner values toward wildlife and wildlife management. Data were collected via a mail survey administered to a randomly-selected sample of landowners with ≥ 160 ac. A total of 1418 landowners participated, resulting in an overall response rate of 47 percent. The majority of landowners indicated that hunting occurs for the species that are present on their lands. Landowners in the eastern region of the state reported the highest levels of hunting occurrence across species, except for elk. The largest difference in hunting occurrence across regions was reported for upland game birds – 63 percent of respondents from the western region with upland game birds on their property allowed access, compared to 80 percent in the East and 74 percent in the central region. For the ungulate species (deer, elk, and antelope), hunting tended to occur more often for male animals. The average number of hunters/year allowed on private lands was highest for elk and lowest for antelope. The two most common forms of access management reported by landowners were “non-Block Management hunting without a fee” involving mostly hunters who are (1) family/friends and (2) NOT family/friends. The third most frequently selected approach was the Block Management Hunting Access Program. By facilitating a better understanding of how hunting is currently managed on private lands in Montana, these results provide a baseline that can assist FWP in developing plans for working with private landowners on issues related to hunting access and wildlife management in the future.

PUBLIC ATTITUDES TOWARDS BLACK BEARS IN MISSOULA, MONTANA

Jerod A. Merkle, Wildlife Biology Program, University of Montana, Missoula, Montana 59801

Paul R. Krausman, Wildlife Biology Program, University of Montana, Missoula, Montana 59801

Melinda M. Booth, Sequoia Park Zoo Foundation and Humboldt State University, Samoa, CA 95564

Successful wildlife management actions and policies depend on public acceptance. Management actions minimizing human-black bear (*Ursus americanus*) conflicts are controversial, and research that articulates public attitudes in a diversity of situations is often

lacking. Our objectives were to examine public attitudes towards black bear management of residents in the Wildland-Urban Interface (WUI; where bears and humans coexist) and the city core, i.e., where no bears are present. We also examined the effects of education and information (EI) efforts to minimize human-bear conflicts. Two questionnaires were distributed to residents living in Missoula, Montana before (2004) and after (2008) EI efforts were implemented. Residents living in the city core and the WUI both attracted bears with bird feeders, BBQ grills, and gardens but had significantly different frequencies of available vegetation and garbage. In the WUI, the frequency of available native and non-native vegetative bear food was significantly higher, whereas the frequency of outdoor garbage storage was significantly lower. In 2008, attitudes were not significantly different between residents in the WUI and the city core. The 4 yrs of EI efforts did not alter resident behaviors that produce attractants in the WUI. However, the EI efforts in the WUI did increase support for non-lethal management actions. Managers may be able to use EI efforts to gain support for black bear management actions, but must realize that behaviors producing attractants may be spatially ubiquitous and difficult to modify.

ASSESSING ECOLOGICAL IMPACTS DUE TO THE OPERATION OF LIBBY DAM, MONTANA

Norm Merz, Fish and Wildlife Division, Kootenai Tribe of Idaho, Bonners Ferry, Idaho 83805

Dwight Bergeron, Montana Fish, Wildlife and Parks, Kalispell, Montana 59901

Damming of rivers represents a cataclysmic event for large river-floodplain ecosystems. By altering water, sediment, and nutrient flow dynamics, dams interrupt and alter a river's important ecological processes in aquatic, riparian, floodplain and surrounding terrestrial environments. The Kootenai Tribe of Idaho and Montana Fish, Wildlife and Parks, with the support of numerous subcontractors, are currently exploring the ecological impacts related to the operation of Libby Dam. To assess the cascading effects, KTOI personnel and subcontractors are developing indices to assess individual abiotic and biotic components. These individual assessments will be combined into an Index of Ecological Integrity (IEI) at a geomorphic reach or subbasin scale. The IEI will be used to assess, mitigation, monitor, and rehabilitate the Kootenai River Floodplain.

LONG-TERM MONITORING OF OSPREY POPULATIONS IN WESTERN MONTANA

Amanda A. Ormesher and Erick Greene, Wildlife Biology Program, University of Montana, Missoula, Montana 59812

Ospreys (*Pandion haliaetus*) are important apex predators and continue to be important indicators for environmental contaminants in aquatic ecosystems. Although osprey populations made rapid recoveries in Montana after the 1972 ban on the use of DDT, there have been surprisingly few studies monitoring osprey populations since then. During the last 20 yrs some osprey populations in western Montana have declined over 75 percent, and are now approaching the low numbers of the DDT era. The causes and geographical extent of these declines are not known, although companion studies have shown some very high levels of mercury in osprey chicks. For these reasons, we are establishing a long-term monitoring program for ospreys in western Montana. Here we report on our osprey studies on Flathead Lake and adjacent watersheds, which appear to be a major stronghold for ospreys. During

the summer of 2008, we located and monitored 87 osprey nests on Flathead Lake. At least 69 (80%) of these nests were occupied; a sub-sample of 47 of these nests had 1.9 chicks produced per productive nest. This is above the productivity estimated to maintain stable osprey populations.

FAT BUT NOT HAPPY: THE EFFECTS OF SUPPLEMENTAL FEEDING ON STRESS HORMONE LEVELS OF WYOMING ELK

Victoria Patrek, Mark Taper, and Scott Creel, Department of Ecology, Montana State University, Bozeman, Montana 59717

Paul Cross, Northern Rocky Mountain Science Center, U.S. Geological Survey, Bozeman, Montana 59717

On 23 feed grounds in Wyoming, elk (*Cervus elaphus*) were provided with supplemental feed throughout the winter. Brucellosis seroprevalence of feed ground elk is 26 percent whereas other elk in the Greater Yellowstone Ecosystem have a brucellosis seroprevalence of 2-3 percent. The aggregation of elk during peak transmission allows brucellosis to persist in the fed populations. In addition to creating the opportunity for disease transmission, the aggregation of elk on feed grounds may have detrimental physiological effects. Other studies have shown that chronically high stress hormone concentrations can suppress the immune system and lead to increased disease susceptibility. Potential stressors on the feed grounds include high densities, large group sizes and aggressive social interactions. In this study we investigated how supplemental feeding and environmental variables affect stress hormone levels in fed and unfed populations of elk. We also manipulated resource distribution on the feed grounds to examine how feeding density affects stress hormone levels and aggression rates. Results show that elk on feed grounds have stress hormone levels significantly higher than unfed elk. Experimental reduction of feed density did not have an effect on stress hormone levels or aggression rates. Group size and density during feeding were important predictors of stress hormone levels. This study indicates that feed grounds are not only creating an epidemiological setting for disease transmission, but also creating a physiological state that may increase susceptibility to disease. The impact of these stress hormone concentrations on disease susceptibility remains unknown, but may potentially be an important driver of disease dynamics in feedground elk populations.

CHANGES IN ELK RESOURCE SELECTION AND DISTRIBUTIONS ASSOCIATED WITH THE MADISON VALLEY LATE-SEASON ELK HUNT

Kelly M. Proffitt and Robert A. Garrott, Department of Ecology, Montana State University, Bozeman, Montana 59717

Changes in resource selection associated with human predation risk may alter elk (*Cervus elaphus*) distributions and availability for harvest. Using Global Positioning System (GPS) data collected from telemetry-collared cow elk, we evaluated effects of refuges (areas where hunting was prohibited), spatial variation in hunting risk and landscape attributes on resource selection within an established Greater Yellowstone Area wintering range and we evaluated elk distributions during and outside of a late-season hunting period. Refuge areas and landscape attributes such as habitat type and snow water equivalency (SWE) affected resource selection. During the hunting period, selection for refuge areas increased, and we estimated odds of elk occupancy in refuge areas more than doubled. Elk selection for flat grasslands

increased as SWE increased likely because these areas are heavily windswept leaving grasses exposed for foraging. Elk distributions differed during hunting and no-hunting periods, and during the hunting period elk distribution shifted to privately owned land where hunting was prohibited. Risk-driven changes in resource selection resulted in distributions that reduced the availability of elk for harvest. Elk selection for areas where hunting is prohibited presents a challenge for resource managers that use hunting as a tool for managing herd sizes.

THE MONTANA LEGACY PROJECT – CONSERVATION OF FOREST LANDS IN WESTERN MONTANA

Robert Rasmussen, The Trust for Public Land, Northern Rockies Field Office, Helena, Montana 59601

The Montana Legacy Project implements an agreement between Plum Creek Timber Company and a partnership between The Nature Conservancy of Montana and The Trust for Public Land for the acquisition of 312,000 ac of Plum Creek lands in western Montana. The goals of the project include: protection of the area's clean water and abundant fish and wildlife habitat, maintenance of forests in productive timber management, and promotion of public access for fishing, hiking hunting and other outdoor recreation. The project includes several areas where Plum Creek lands are in a checkerboard ownership pattern with public lands. The potential conversion of these lands to subdivision and development could significantly fragment wildlife habitat, interrupt movement corridors, restrict traditional public recreational access, and increase the costs of wildfire protection and the provision of local services. This project is the result of collaboration by many individuals and organizations from across the state. Business leaders, sportsmen, elected officials, federal and state agencies, conservationists, citizen groups and others are working together to conserve the important resource values within this project area.

ESTIMATION OF BLACK-TAILED PRAIRIE DOG COLONY ACREAGE IN MONTANA

Ryan Rauscher, Montana Fish, Wildlife & Parks, Glasgow, Montana 59230

Scott Story, Montana Fish, Wildlife & Parks, Helena, Montana 59620

Justin Gude, Montana Fish, Wildlife & Parks, Helena Montana 59620

The first statewide inventory of black-tailed prairie dog (*Cynomys ludovicianus*) colonies in Montana was conducted in the mid 1980s and produced an estimate of 120,000 – 130,000 ac. A cooperative statewide inventory effort (1996 -1998) yielded a minimum estimate of 66,000 ac of occupied habitat. However, this inventory did not include areas where access was denied and results of alternate survey efforts. The minimum management objective for black-tailed prairie dog abundance in Montana is 90,000–104,000 occupied ac, according to the statewide management plan. A current, comprehensive, and rigorous statewide estimate of prairie dog acreage was needed to determine the area of Montana occupied by black-tailed prairie dog colonies. We surveyed known black-tailed prairie dog distribution in Montana using an aerial line-intercept survey stratified by county from June to August, 2008. We recorded the length of prairie dog colony intercepts on approximately 35,000 linear mi on 771 transects in 32 counties in central and eastern Montana. Transects were flown in fixed-wing aircraft in an east-west direction at ~ 110 mph at < 300 ft above ground level. Prairie dog

colonies were detected in 21 counties. We recorded roughly 750 prairie dog colony intercepts. We attempted to ground truth 10 percent of detected prairie dog colonies along transects, and at least one black-tailed prairie dog colony intercept in each county. Data analysis is ongoing at this time. However, preliminary results indicate black-tailed prairie dog occupancy exceeds Montana abundance objectives. A comprehensive analysis is expected to be complete in April, 2009.

EFFECTS OF CATTLE GRAZING ON SMALL MAMMAL COMMUNITIES AT RED ROCK LAKES NATIONAL WILDLIFE REFUGE

Greg Reed, Department of Ecology, Montana State University, Bozeman, Montana 59717

Nathan Whelham, Department of Intercollege Programs for Science Education, Montana State University, Bozeman, Montana 59717-2805

Jeffrey M. Warren, USDI Fish and Wildlife Service, Red Rock Lakes National Wildlife Refuge
Lima, Montana 59739

Michael R. Frisina, Montana Fish, Wildlife and Parks, Butte, Montana 59701

Cattle grazing is a common land practice on public lands in the West that can have complex impacts on both wildlife and vegetation. However, many studies of wildlife response to grazing only compare grazed versus ungrazed treatments, ignoring the dynamic nature of grazing and the many levels of grazing intensity and frequency commonly utilized. We undertook the current study to better understand the response of small mammals to the frequency of cattle grazing in wet meadow habitats on Red Rock Lakes NWR. Three adjacent grazing units were selected for study with grazing frequencies of 1, 3, and 8 years of rest in 2007 and 0, 2, and 4 yrs of rest in 2008. Two randomly placed trapping grids were placed within the *Juncus balticus*–*Carex praegracilis* vegetative alliance (wet meadow) in each unit. Trapping occurred from late June–August each year. Vegetation was quantified in each unit each year using point-intercept transects. We conducted raptor surveys during 2008 to examine raptor response to vole abundance. Voles (meadow and montane; *Microtus* spp.) made up the overwhelming majority (~99%) of individuals captured. Results from 2007 indicated that vole abundance increased with increasing rest from grazing. Vole abundance in 2008 was substantially lower and did not follow the same pattern as 2007—in 2008 the unit with an intermediate level of rest (2 yrs) had the highest abundance. Raptor numbers tracked vole abundance closely in 2008. Additional years of study will be necessary to determine the role grazing plays in vole population dynamics.

RECENT ADVANCES IN THE ANALYSIS OF OCCUPANCY AND ABUNDANCE DATA IN RESPONSE TO MANAGEMENT ACTIVITIES

Robin Russell, Montana Fish, Wildlife, and Parks, Bozeman, Montana 59718

Recent advances in the analysis of occupancy data collected from repeated visits to sampling stations will be presented along with an example analysis of avian community responses to prescribed fire. The occupancy analysis is a Bayesian hierarchical model, allows fixed and random effects, and is a composite analysis that allows researchers to estimate management effects for rare species as well as community level indices such as species richness. Additional examples of applied research projects that have used Bayesian hierarchical models will be briefly discussed to demonstrate the flexibility of the method and

introduce the audience to techniques they may not be familiar with. Additionally, an example using variable circular plot, i.e., distance data, will be presented demonstrating an analysis of red squirrel (*Tamiasciurus hudsonicus*) densities in response to prescribed fire. These methods are appropriate for studies designed to investigate the impact of management practices on wildlife species and represent improvements over previous techniques.

AS THE WORLD TURNS: MANAGING WOLVES AND NAVIGATING THE LEGAL LABYRINTH

Carolyn A Sime, Elizabeth Bradley, Kent Laudon, Mike Ross, and Nathan Lance, Montana Fish, Wildlife and Parks, Helena, Montana 59620

Val Asher, Turner Endangered Species Fund, Bozeman, Montana 59718

This is the 29th year that wolves (*Canis lupus*) have been in northwest Montana and the 14th year that wolves have been in southwest Montana. It's the 5th year Montana Fish, Wildlife and Parks (FWP) has led wolf management. And it's the 7th straight year in which the northern Rockies biological recovery goal was met. The real success story is that roughly 88 percent of the recovered wolf population lives outside national parks. Through it all, science has advanced and the rhetoric has ebbed and flowed. Somewhere in the midst are the biological, political, and social realities of managing a recovered population. Outside national parks, wolves share a landscape with people who have very diverse viewpoints about wolves and their management. Like other wildlife species, Montana's wolf population is subject to checks and balances, including strong reproduction in some areas, disease, vehicle strikes, and mortality due to conflicts with people. Montana's population is secure, but dynamic. At the end of 2007, Montana had a minimum of 422 wolves in 73 packs, 39 of which were breeding pairs. In 2008, the legal framework was equally dynamic when federal government efforts to delist the northern Rockies population were challenged. Wolves were delisted for about 4 mos before a federal judge reinstated federal legal protections through a preliminary injunction and then permanently when the federal government withdrew its decision. Delisting efforts were renewed late in 2008 and carried forward into 2009. This will be summarized and an FWP program update will be provided. See also: www.fwp.mt.gov/wildthings/wolf

CCAA IMPACTS ON BIRDS OF THE BIG HOLE WATERSHED

Kristina Smucker and Megan Fylling, Avian Science Center, Health Science 209, Division of Biological Sciences, University of Montana, Missoula, Montana 59812

Amy Cilimburg, Montana Audubon, Helena, Montana 59624

Through the Big Hole CCAA, a coalition of groups has undertaken extensive restoration work aimed at improving habitat conditions for the last remaining fluvial population of Arctic grayling in the U.S. Riparian-associated birds are also likely to respond positively to such restoration activities, and this is significant because riparian habitats support a greater diversity of breeding birds than any other habitat type in Montana. In the summers of 2007 and 2008 the Avian Science Center obtained permission from participating landowners along the Big Hole River to document bird communities during this pre-restoration phase. We surveyed birds at reference points that serve as the target habitat condition for restoration efforts, and impact points that are currently impaired, but likely to improve with restoration activities. We also measured vegetation at each point to evaluate vegetation quality. We

detected 107 species across the two survey years, and this represents 44 percent of bird species (107/245) known to breed in Montana. This speaks to the outstanding diversity of birds associated with riparian areas in the Big Hole watershed. Both the vegetation characteristics and bird communities were decidedly different between reference and impact sites. Thirteen of the 25 most frequently encountered bird species were detected on a significantly higher proportion of reference points while six species were more frequently encountered on impact sites. We suggest using the occurrence of five species: willow flycatcher (*Empidonax trailii*), veery (*Catharus fuscescens*), northern waterthrush (*Seiurus noveboracensis*), fox sparrow (*Passrella iliaca*) and song sparrow (*Melospiza melodia*), all of which were encountered much more frequently at reference sites, to evaluate the success of restoration work in future years.

TE IPOR 1. COMPARISONS OF GREAT BLUE HERON ROOKERY DISTRIBUTION, ABUNDANCE AND REPRODUCTIVE SUCCESS IN THE LOWER YELLOWSTONE RIVER BASIN

Dean J. Waltee,* Montana Fish, Wildlife & Parks, Miles City, Montana 59301

Ryan L. Rauscher, Montana Fish, Wildlife & Parks, Glasgow, Montana 59230

In response to declining trend information from Montana Audubon Society, Breeding Bird Surveys, and as part of an ongoing effort to assess the status of eastern Montana's non-game wildlife and identify critical habitats, we assessed the distribution and abundance, occupancy and reproductive success of great blue heron (*Ardea herodias*) rookeries within the lower Yellowstone River Basin. We conducted an aerial census of the lower Yellowstone, Tongue and Powder river corridors to document existing rookeries. Rookeries were revisited prior to fledgling abandonment to assess reproductive success. We compared the distribution, abundance and reproductive success of rookeries along the lower Yellowstone River to data from similar surveys conducted in 1976 and 1988. Great blue heron rookery abundance and reproductive success were also compared to river flows during corresponding time periods. Between 1976 and 1988, active rookery abundance increased from nine to 18 but declined back to nine by 2008. The average number of occupied nests per active rookery for each time period was 14 in 1976, 16.8 in 1988 and 12.1 in 2008. Rookery distribution showed temporal variance, however, some areas supported rookeries during all survey periods. Reproductive success data were not available for 1976. Comparisons between 1988 and 2008 showed that the percentage of reproductively successful rookeries in 1988, when river flows remain primarily below average, was 61 percent. During 2008, when river flows were consistently above average, the percentage of successful rookeries decreased to 44 percent. Combined data suggested that great blue heron rookery abundance and reproductive success along the lower Yellowstone River fluctuates temporally in response to various ecological factors, yet to be fully understood.

MANAGING BLACK BEARS AND COUGARS WITH PEOPLE PROBLEMS

Erik Wenum, Montana Fish, Wildlife and Parks, Kalispell, Montana 59901

The Flathead Valley area and most of northwest Montana is currently growing at a rate of 6 percent/ yr. Given the increasing human population and expanding residential development in the urban interface, there will continue to be bears (*Ursus* spp.) and cougars

(*Puma concolor*) living in close proximity to homes and human activity centers. During the 2007 and 2008 field seasons we received 774 and 881 black bear (*Ursus americanus*) conflict calls, respectively, concerning incidents of some kind. Most of these black bear calls occurred between May 1 and November 30, during 2008 that results in 881 calls in a 7-mo period or 126 calls/mo or 4.1 calls/day – everyday. During the 2007 and 2008 field seasons we received 70 and 152 cougar conflict calls, respectively, concerning cougar conflicts of some kind. The 6 yrs previous to 2008 we received an average of 68 calls/year concerning cougar conflicts. This dramatic increase in cougar conflicts in 2008 reflects an increasing cougar population trend in the last few years. Through continued education efforts it will be possible to disseminate the best available information to pre-emptively reduce human / wildlife conflicts. The continued use of a bear trap to relocate or trap and aversively condition program will address bears that have already made a positive association with people or their dwellings. The continued response to cougar conflicts will reduce (though not eliminate) the potential public safety issues that exist anywhere there are healthy cougar populations. It is our hope that we will be able to maintain the public tolerance for these highly prized big game animals in western Montana by maintaining this effective wildlife conflict, safety and education program.

BIRD DISTRIBUTION IN MONTANA: OPPORTUNITIES FOR PUBLIC INVOLVEMENT

Catherine Wightman and Scott Story, Montana Fish, Wildlife and Parks, Helena, Montana 59620

Coburn Currier, Montana State Library, Helena, Montana 59620

Montana's diverse landscapes provide habitat to a wide assortment of birds. Climate, ecology, and human activity can influence landscape changes and patterns of bird distributions across the state. Every year, birdwatchers, biologists, and landowners document the occurrence of common and rare bird species throughout Montana. The Montana Bird Distribution program promotes stewardship of birds and their habitats by using information gathered by Montana's citizens to track the distribution and seasonal occurrence of the 409 bird species documented in the state. Over 30 new species have been documented in the state of Montana since 1980 through this program. As of 2003, 29 species appeared to be increasing their distribution across the state, e.g., tundra swans, several flycatcher species, and 27 species appeared to be decreasing their distribution, e.g., stilt sandpipers, 2 species of rosefinch. The Montana Bird Distribution partnership consists of Montana Audubon, Montana Bird Records Committee, Montana Natural Heritage Program, and Montana Fish, Wildlife and Parks. The Partnership collects observations by professional and nonprofessionals, maintains a high-quality database of observations, and disseminates information via a website and *P.D. Skaar's Montana Bird Distribution* book series. Publication of the next book edition is planned for 2010 and observations are due by 31 December 2009 for inclusion in this edition. Here we present examples from the previous edition and a tutorial on *Tracker*, the web-based system for contributing data points.

PATTERNS OF MOVEMENT IN BLACK-BACKED WOODPECKERS

Jennifer C. Woolf, Wildlife Biology Program, University of Montana, Missoula, Montana 59812

Fred W. Allendorf, Division of Biological Sciences, University of Montana, Missoula, Montana 59812

Michael Schwartz, USDA Forest Service, Rocky Mountain Research Station, Missoula, Montana 59801

Black-backed woodpeckers (*Picoides arcticus*) inhabit recently burned forests, forcing both sexes to regularly disperse during the course of their lifetime. Yet we know little about their pattern of dispersal, including both distance traveled and the range of habitat types colonized for reproduction. We collected genetic samples from 274 black-backed woodpeckers across North America. We found a large amount of gene flow across the boreal forest and used a simulation approach to test if this pattern is best explained by frequent colonization of burned patches. We then used the fire history in the boreal forest to estimate the average distance between fires to determine if burned patches are readily available for colonization. Finally, we estimated the average dispersal distance based on the genetic relatedness among birds in our study. In combination, this information can be used to help prioritize land management decisions, such as salvage logging, that affect black-backed woodpecker populations.

A GIS TOOL FOR CONDUCTING LANDSCAPE-SCALE HABITAT QUALITY ASSESSMENTS

Scott D. Yeats, Jonathan B. Haufler, and Carolyn A. Mehl, Ecosystem Management Research Institute, Seeley Lake, Montana 59868

Evaluating and quantifying human impacts and improvements to wildlife habitat at large scales is a continuing challenge for wildlife managers. To help such evaluations, we have developed a GIS-based habitat modeling tool that quantifies the number and quality of home ranges for species of interest in existing, historical, or potential future landscape conditions. We used this tool in a 1.5-million ac landscape in northern Idaho to model habitat-based species viability for eight species of concern. This presentation will focus on the results for Canada lynx, northern goshawk, and flammulated owl. We developed habitat potential maps at the scale of the home range for each species. Models were created for existing, historical, and predicted future habitat conditions. Future habitat conditions were projected based on an objective of maintaining or restoring 20 percent representation of historical conditions. Model runs were used to produce an estimate of the number of high, moderate, low, and very low quality home ranges, with implications from this for likely persistence of each species. Improvements in future habitat quality varied among the species, but improved the most for those species dependent upon drier forest types where greater restoration efforts are needed and planned. The ability to model habitat-based species viability allows land managers to estimate the impacts or improvements to wildlife species of concern that are likely to result from development, management or restoration activities.

TERRESTRIAL POSTER SESSION

MODELING PREDICTED DISTRIBUTION AND LANDSCAPE LEVEL HABITAT SUITABILITY FOR MONTANA WILDLIFE SPECIES

Bryce A. Maxell, Montana Natural Heritage Program, University of Montana, Helena, Montana 59620

Scott Story and Joy Ritter, Information Management Bureau, Montana Department of Fish, Wildlife, and Parks, Helena, Montana 59620

Models predicting spatial distribution and habitat suitability are critical for natural resource managers making decisions that impact species for which there is limited information. We are using presence-only data in conjunction with pseudo-absences in program Maxent to model distribution and landscape-level habitat suitability for Montana wildlife species. Our primary goals are to produce (1) continuous statewide outputs as a tool to identify variables that limit species' distributions and areas that need field surveys; (2) binary outputs that can be used to create lists of predicted species for various administrative boundaries and, (3) outputs showing marginal, suitable, and optimal habitat classes at a local landscape-level. To date, models have identified scale-dependent responses to environmental variables, opportunities to extend the known ranges of species, areas that support potentially isolated populations in need of conservation efforts, areas that are critical for maintaining landscape connectivity, areas that may provide the best habitat for reintroduction of species that have declined, and areas where exotic and nonindigenous species are most likely to become established. In general, inductively based Maxent models provide realistic depictions of species distributions when survey data is available for a region. However, deductive models will still be important for representing some species distributions in areas lacking survey effort. Models will be used for various planning efforts including Montana Fish, Wildlife, and Parks' Crucial Areas and Corridors Assessment. Model outputs can be obtained from the Montana Natural Heritage Program or the Montana Fish, Wildlife, and Parks' Information Management Bureau.

STATUS OF LENTIC BREEDING AMPHIBIANS AND AQUATIC REPTILES IN MONTANA

Bryce A. Maxell and Dave Ratz, Montana Natural Heritage Program, University of Montana, Helena, Montana 59620

P. Stephen Corn, U.S. Geological Survey Northern Rocky Mountain Science Center, Aldo Leopold Wilderness Research Institute, Missoula, Montana 59807

D. Grant Hokit, Department of Natural Sciences, Carroll College, Helena, Montana 59625

We developed a statewide inventory and monitoring scheme for lentic breeding amphibians and aquatic reptiles in Montana. We stratified sampling by 11 ecoregions and surveyed 6741 potential lentic sites on public lands between 2000 and 2008 within 429 randomly selected 12-digit hydrologic-unit-code watersheds. Surveys and associated incidental observations have resulted in over 11,400 species occurrence records to date. Watershed and site breeding rates of 10 amphibian species and occupancy rates of four aquatic reptile species support previously noted declines of the Western Toad (*Bufo boreas*)

and Northern Leopard Frog (*Rana pipiens*) in western Montana. We used classification trees to examine patterns in rates resulting from different groupings of major habitat features that are able to be affected with management actions. Seven of the 10 amphibian species and the Common Gartersnake (*T. sirtalis*) were detected at significantly fewer sites when fish were detected. Presence of emergent vegetation was positively associated with the proportion of sites where breeding or occupancy was detected for all but one of the amphibian and reptile species examined and appeared to partially mitigate the presence of fish. Resource managers could enhance habitats for wetland herpetofauna by (1) creating new lentic sites on the landscape either directly or through the reintroduction and protection of beaver, (2) creating emergent vegetation at portions of existing sites that currently lack it via rotational fencing to temporarily exclude grazing, and (3) eliminating some introduced fish populations. All observations and survey locations, including digital photographs of sites surveyed are available at: <http://hhp.nris.state.mt.us/Tracker>.

THE PATHOGENIC CHYTRID FUNGUS IN MONTANA AMPHIBIANS

Bryce A. Maxell, Montana Natural Heritage Program, University of Montana, Helena, Montana 59620

Blake R. Hossack and P. Stephen Corn, U.S. Geological Survey Northern Rocky Mountain Science Center, Aldo Leopold Wilderness Research Institute, Missoula, Montana 59807

Erin Muths, U.S. Geological Survey Fort Collins Science Center, Fort Collins, Colorado 80525

David S. Pilliod, U.S. Geological Survey Forest and Rangeland Ecosystem Science Center, Snake River Field Station, Boise, Idaho 83706

Janene Lichtenberg, Tribal Wildlife Management Program, Confederated Salish and Kootenai Tribes, Pablo, Montana 59855

J. Kirwin Werner, Department of Environmental Sciences, Salish Kootenai College, P.O. Box 70, Pablo, Montana, 59855

Carter Kruse, Turner Enterprises, Inc., Bozeman, Montana 59718

D. Grant Hokit, Department of Natural Sciences, Carroll College, Helena, Montana 59625

D. Earl Green, U.S. Geological Survey National Wildlife Health Center, Madison, Wisconsin 53711-6223

The chytrid fungus (*Batrachochytrium dendrobatidis*; Bd) is pathogenic to many amphibians and has been linked to declines and extinctions in a number of species around the globe. We collected 484 tissue samples and swabs of ventral surfaces of nine species at 161 locations across Montana in the course of fieldwork between 1998 and 2005. We detected Bd in 161 samples taken at 64 sites between 1998 and 2005 for six species; *A. tigrinum*, *B. boreas*, *B. woodhousii*, *P. maculata*, *R. luteiventris*, and *R. pipiens*. Overall, 40 percent of sites had at least one species test positive and, across all species, 33 percent of samples tested positive. We detected Bd in samples taken throughout the active season of these species across Montana at a variety of elevations, in a variety of habitats, and up to 17.5 km from the nearest road. However, a higher percentage of samples and sites (37 to 41 and 47 to 66 percent, respectively) tested positive for Bd within 1 km of roads. While this effort provides no definitive evidence, Bd, acting alone or synergistically with other stressors, appears to be the most likely cause of declines observed in *B. boreas* and *R. pipiens* populations in western Montana in light of its widespread distribution and association with declines in other regions. Further evaluation of the status of Bd and other amphibian pathogens in Montana is warranted and strict adherence to protocols preventing the spread of these pathogens is needed.

IMPROVING SCIENCE LITERACY THROUGH VISUAL COMMUNICATION

Mary McFadzen and Melissa Brown, Center for Invasive Plant Management, Land Resources and Environmental Sciences Department, Montana State University, Bozeman, Montana 59717

Communicating science effectively to diverse audiences including natural resource managers, policy makers, and the general public, has broad implications for implementing science-based solutions to environmental problems such as those associated with non-native species invasions. If audiences are able to understand and assess scientific information, they are more likely to make decisions based on facts rather than on hearsay or speculation. Frequently, popular press articles on invasive species are designed to promote fear-based responses. Previous work in an ecological publication identified that even invasion biologists use militaristic and combative language as metaphors, often leading to an inaccurate perception of invasive species and loss of scientific credibility, which can be counterproductive to achieving conservation and management goals. Scientists and educators are natural collaborators to increase the science literacy about the biology and ecology of non-native species invasions to audiences, while consistently maintaining scientific integrity and credibility. Further, as increasing numbers of research granting entities require an outreach and education component in project proposals and outputs, proven ability to effectively present and disseminate scientific information to diverse audiences is becoming highly valued. This poster presents examples of how visual design strategies and learning theory can be used to engage audiences and communicate science effectively.

SEROPREVALENCE OF CANINE PARVOVIRUS AND CANINE DISTEMPER IN WOLVES IN RELATION TO HUMAN ACTIVITY IN THE CANADIAN ROCKY MOUNTAINS

Brynn Nelson, Wildlife Biology Program, University of Montana, Bonner, Montana 59823

Mark Hebblewhite, Wildlife Biology Department, University of Montana, Missoula, Montana 59802

Todd Shury, Parks Canada, Department of Veterinary Pathology, Western College of Veterinary Medicine, Saskatoon, Saskatchewan, Canada S7N 5B4

Evelyn Merrill, Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2E9

Dale Seip, BC Ministry of Forests, Prince George, British Columbia, Canada V2L3H9

Nathan Webb, Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2E9

Fiona Schmiegelow, Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada T6G 2H1

Paul C. Paquet, World Wildlife Fund, Meacham, SK, Canada S0K 1V0

Diseases affect social carnivores that occur in high density areas, like wolves (*Canis lupus*). Carrier species (feral dogs, coyotes, foxes) travel between the urban/wildlife interface; thus, transmitting diseases to wolves. We sampled 99 wolves from the years 2000 to 2008 for canine parvovirus (CPV) and canine distemper virus (CDV) in Banff and Jasper National Parks and surrounding areas of the Canadian Rockies. Of the 99 wolves, 92 tested positive

for CPV, 22 tested positive for CDV and 22 tested positive for both diseases. We tested whether seroprevalence of CPV and CDV was higher closer to human activity (roads, town sites, campgrounds, federally designated Indian reserves) and as a function of sex, age class, and different wolf packs using mixed-effects logistic regression models. CPV and CDV seroprevalence was found to be higher in areas closer to human activity and was higher in younger age classes of wolves. Understanding disease transmission between urban areas and wildlife areas with high wolf densities, like the Canadian Rockies, could yield pertinent information about disease profiles. Disease profiles from the Canadian Rockies could help conserve the recently delisted wolf species in areas like Yellowstone National Park where human activity is high relative to wolf activity.

DYNAMICS OF A HARVESTED MOOSE POPULATION IN SOUTHWEST MONTANA

Megan O'Reilly and Jeffrey M. Warren, USDI Fish and Wildlife Service, Red Rock Lakes National Wildlife Refuge, Lima, Montana 59739

Bob Brannon, Montana Fish, Wildlife and Parks, Sheridan, Montana 59749

Floodplain riparian habitat provides relatively stable and important wintering habitat for moose in the Rocky Mountains. The Centennial Valley (CV) in southwest Montana contains the largest wetland complex in the Greater Yellowstone Ecosystem, and the associated riparian habitat supports one of the largest and highest-density wintering Shiras moose (*Alces alces shirasi*) populations in the Northern Rocky Mountains. Most of this habitat is encompassed by Red Rock Lakes National Wildlife Refuge (Refuge), situated in the eastern extent of the valley and within Montana Fish, Wildlife and Parks hunting district 334 (HD 334). We utilized 4 decades of winter aerial survey (1966-2008) data from HD 334 to investigate relationships among winter moose abundance, snow pack, and hunter harvest of antlerless moose. Winter moose abundance steadily increased during the period 1966-2008 at an average annual rate of 2.2 percent (SE = 0.48%), but per capita productivity (calf: adult ratio) simultaneously declined. As predicted, winter snow pack was positively related to moose abundance on the survey area. Hunter harvest of antlerless moose negatively affected moose abundance, but the effect differed for the two age classes (calves vs. adults). Our results indicate that previous levels of hunter harvest were sustainable; however this occurred at a time when few other moose predators were present. Future harvest management will have to take into account increasing numbers of predators in southwestern Montana.

SPONSORING ORGANIZATIONS AND 2009 OFFICERS

THE MONTANA CHAPTER OF THE AMERICAN FISHERIES SOCIETY

President	Scott Barndt USFS Gallatin National Forest - Bozeman, MT
President Elect	Todd Koel NPS Yellowstone National Park - WY
Past President	Carter Kruse Turner Enterprise, Inc. - Bozeman, MT
Secretary-Treasurer	Leo Rosenthal Montana Fish, Wildlife and Parks - Kalispell

THE MONTANA CHAPTER OF THE WILDLIFE SOCIETY

President	Joe Weigand Montana Fish, Wildlife and Parks - Helena
President Elect	Julie Cunningham Montana Fish, Wildlife and Parks - Bozeman
Past President	Ryan Rauscher Montana Fish, Wildlife and Parks - Glasgow
Secretary	Jay Kolbe Montana Fish, Wildlife and Parks - Selley Lake
Treasurer	Mike McGrath Montana Department of Natural Resources & Conservation - Missoula

MONTANA ACADEMY OF SCIENCES

President	James Barron Montana State University - Billings
Past President	Doug Coffin University of Montana - Missoula
Executive Director	Doug Coffin University of Montana - Missoula
Recording Secretary	Carmen Hauck Big Sky High School - Missoula, MT
Asst. Recording Sec.	Laurie Henneman University of Montana -Dillon
Treasurer	Tom Lewis Montana State University - Billings

Intermountain Journal of Sciences

Vol. 15, No. 4 - 2009

CONTENTS

ARTICLES

Biological Sciences - Aquatic

New Zealand Mudsail Containment: Tests of Chemicals, Copper Strips,
Fluorochrome Marking and Terrestrial Movement41
Randall W. Oplinger and Eric J. Wagner

Health and Human Development

Effects of a 6-week Jump Training Program Combined with Two Lifting
Programs54
John Amtmann, Jeremy Basler, Charles S. Todd and Shawn Vincent

ABSTRACTS

Plenary Session58
Biological Sciences - Aquatic65
Biological Sciences - Terrestrial90